

Product datasheet for **SC329685**

IL17RC (NM_001203264) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	IL17RC (NM_001203264) Human Untagged Clone
Tag:	Tag Free
Symbol:	IL17RC
Synonyms:	CANDF9; IL17-RL; IL17RL
Vector:	pCMV6-Entry (PS100001)




[View online »](#)

Fully Sequenced ORF: >SC329685 representing NM_001203264.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

ATGCCTGTGCCCTGGTTCTTGCTGTCTTGGCACTGGGCCGAAGCCAGTGGTCTTTCTCTGGAGAGG
CTTGTGGGGCCTCAGGACGCTACCCACTGCTCTCCGGGCCTCTCCTGCCGCCTCTGGGACAGTGACATA
CTCTGCCTGCCTGGGGACATCGTGCCTGCTCCGGGCCCGTGTGGCCCTACGCACCTGCAGACAGAG
CTGGTGTGAGGTGCCAGAAGGAGACCGACTGTGACCTGTGTCTGCGTGTGGCTGTCCACTTGGCCGTG
CATGGGCACTGGGAAGAGCCTGAAGATGAGGAAAAGTTTGGAGGAGCAGCTGACTCAGGGGTGGAGGAG
CCTAGGAATGCCTCTCTCCAGGCCAAGTCGTGCTCTCCTTCCAGGCCTACCCTACTGCCCGCTGCGTC
CTGCTGGAGGTGCAAGTGCCTGCTGCCCTTGTGCAGTTTGGTCAGTCTGTGGGCTCTGTGGTATATGAC
TGCTTCGAGGCTGCCCTAGGGAGTGAGGTACGAATCTGGTCTATACTCAGCCCAGGTACGAGAAGGAA
CTCAACCACACACAGCAGCTGCCTGACTGCAGGGGGCTCGAAGTCTGGAACAGCATCCCAGCTGCTGG
GCCCTGCCCTGGCTCAACGTGTCAGCAGATGGTGACAACGTGCATCTGGTTCTGAATGTCTCTGAGGAG
CAGCACTTCGGCCTCTCCCTGTACTGGAATCAGGTCCAGGGCCCCAAAACCCCGTGGCAGAAAAAC
CTGACTGGACCGCAGATCATTACCTTGAACCACACAGACCTGGTTCCTGCTCTGTATTAGGTGTGG
CCTCTGGAACCTGACTCCGTTAGGACGAACATCTGCCCTTCCAGGGAGGACCCCGCGCACACCAGAAC
CTCTGGCAAGCCGCCGACTGCGACTGCTGACCCTGCAGAGCTGGCTGCTGGACGCACCGTGTCTCGCTG
CCCGCAGAAGCGGCACTGTGCTGGCGGGCTCCGGGTGGGGACCCCTGCCAGCCACTGGTCCCACCCTT
TCCTGGGAGAAGTCACTGTGGACGTGAACAGCTCGGAGAAGTGCAGCTGCAGGAGTGTGTGGGCT
GACTCCCTGGGGCCTCTCAAAGACGATGTGCTACTGTTGGAGACACGAGGCCCCAGGACAACAGATCC
CTCTGTGCCTTGGAAACCCAGTGGCTGTACTTCACTACCAGCAAAGCCTCCACGAGGGCAGCTCGCCTT
GGAGAGTACTTACTACAAGACCTGCAGTCAGGCCAGTGTCTGCAGCTATGGGACGATGACTTGGGAGCG
CTATGGGCCTGCCCATGGACAAATACATCCACAAGCGTGGGCCCTCGTGTGGCTGGCCTGCCTACTC
TTTGCCGCTGCGCTTCCCTCATCCTCCTCTCAAAAAGGATCACGCGAAAGCGGCCGCCAGGGGGCCGC
GCGGCTCTGCTCCTACTCAGCCGATGACTCGGGTTTCGAGCGCCTGGTGGGCGCCTGGCGTCCGCC
CTGTGCCAGTGCCGCTGCGCGTGGCCGTAGACCTGTGGAGCCGTGCTGAACTGAGCGCGCAGGGGCC
GTGGCTTGGTTTACGCGCAGCGCGCCAGACCCTGCAGGAGGGCGGCGTGGTGGTCTTGTCTTCTCT
CCCGGTGCGGTGGCGCTGTGCAGCGAGTGGCTACAGGATGGGGTGTCCGGGCCCGGGGCGCACGGCCG
CACGACGCCTCCGCGCCTCGCTCAGCTGCGTGTGCCGACTTCTTGCAGGGCCGGGCGCCCGGCAGC
TACGTGGGGCCTGCTTCGACAGGCTGCTCCACCCGACGCGTACCCGCCCTTTCCGACCCGTGCC
GTCTTCACTGCCCTCCAACTGCCAGACTTCTGGGGCCCTGCAGCAGCCTCGCGCCCGCGTTC
GGGCGGCTCCAAGAGAGAGCGGAGCAAGTGTCCCGGGCCCTTCCAGCCAGCCCTGGATAGCTACTTCAT
CCCCGGGGACTCCCGCGCCGGGACGCGGGGTGGGACCAGGCGCGGGACCTGGGGCGGGGACGGGACT
TAA
  
```

Restriction Sites: Sgfl-Mlul

Plasmid Map:


ACCN: NM_001203264

Insert Size: 2073 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001203264.1](#)

RefSeq Size: 2362 bp

RefSeq ORF: 2073 bp

Locus ID: 84818

UniProt ID: [Q8NAC3](#)

Cytogenetics: 3p25.3-p24.1

Protein Families: Druggable Genome, Transmembrane

MW: 75.3 kDa

Gene Summary:

This gene encodes a single-pass type I membrane protein that shares similarity with the interleukin-17 receptor (IL-17RA). Unlike IL-17RA, which is predominantly expressed in hemopoietic cells, and binds with high affinity to only IL-17A, this protein is expressed in nonhemopoietic tissues, and binds both IL-17A and IL-17F with similar affinities. The proinflammatory cytokines, IL-17A and IL-17F, have been implicated in the progression of inflammatory and autoimmune diseases. Multiple alternatively spliced transcript variants encoding different isoforms have been detected for this gene, and it has been proposed that soluble, secreted proteins lacking transmembrane and intracellular domains may function as extracellular antagonists to cytokine signaling. [provided by RefSeq, Feb 2011]

Transcript Variant: This variant (5) lacks 2 in-frame coding exons (exon 12 and penultimate exon) compared to variant 1, resulting in a shorter isoform (5, also known as IL-17RCdelta12) missing 2 internal protein segments compared to isoform 1. It has been reported that isoforms lacking exon 12 lose the ability to bind cytokines, IL-17A and IL-17F (PMID:17911633).