

Product datasheet for SC329720

CHODL (NM 001204176) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: CHODL (NM_001204176) Human Untagged Clone

Tag: Tag Free
Symbol: CHODL

Synonyms: C21orf68; MT75; PRED12

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC329720 representing NM_001204176.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GAAGTATAA

Restriction Sites: Sgfl-Mlul


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001204176

Insert Size: 699 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001204176.1</u>

 RefSeq Size:
 2316 bp

 RefSeq ORF:
 699 bp

 Locus ID:
 140578

 UniProt ID:
 Q9H9P2

 Cytogenetics:
 21q21.1

Protein Families: Transmembrane

MW: 26 kDa

Gene Summary: This gene encodes a type I membrane protein with a carbohydrate recognition domain

characteristic of C-type lectins in its extracellular portion. In other proteins, this domain is involved in endocytosis of glycoproteins and exogenous sugar-bearing pathogens. This protein localizes predominantly to the perinuclear region. Several transcript variants encoding a few different isoforms have been found for this gene. [provided by RefSeq, Feb

2011]

Transcript Variant: This variant (4) differs in the 5' UTR and coding sequence compared to variant 1. The resulting isoform (c) is shorter at the N-terminus compared to isoform a.

Variants 3 and 4 both encode the same isoform (c).