

## Product datasheet for **SC329807**

### ATP5MD (NM\_001206426) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	ATP5MD (NM_001206426) Human Untagged Clone
Tag:	Tag Free
Symbol:	ATP5MD
Synonyms:	bA792D24.4; DAPIT; HCVFTP2; MC5DN6; USMG5
Mammalian Cell Selection:	None
Vector:	<u><a href="#">pCMV6-XL5</a></u>
E. coli Selection:	Ampicillin (100 ug/mL)
Fully Sequenced ORF:	>OriGene ORF within SC103736 sequence for NM_032747 edited (data generated by NextGen Sequencing) ATGGCAGGKCCAGAAAAGTGATGCGCAATACCAGTTCCTGTTATTAATAATTTCAAC TCTTATACTCTCACAGGTAGAATGAACTGTGACTGGCCACATATGGAAGCATTGCATTG ATTGTCTTATTTCAAGTTAAGGTCCAAAAAACTCCAGCTGTGAAAGCAACATAA  Clone variation with respect to NM_032747.3 9 t=>k
Restriction Sites:	Sgfl-MluI
ACCN:	NM_001206426
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:custsupport@origene.com">custsupport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)



[View online »](#)

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u>NM_001206426.1</u> , <u>NP_001193355.1</u>
<b>RefSeq Size:</b>	427 bp
<b>RefSeq ORF:</b>	177 bp
<b>Locus ID:</b>	84833
<b>UniProt ID:</b>	<u>Q96IX5</u>
<b>Cytogenetics:</b>	10q24.33
<b>Protein Families:</b>	Transmembrane
<b>Gene Summary:</b>	<p>Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation (Probable). Minor subunit required to maintain the ATP synthase population in the mitochondria (PubMed:21345788). [UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (1) is a predominant transcript. Variants 1, 2 and 3 encode the same protein.</p>