

## **Product datasheet for SC329812**

## ATF3 (NM 001206486) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

Product Name: ATF3 (NM\_001206486) Human Untagged Clone

Tag: Tag Free Symbol: ATF3

**Vector:** pCMV6-Entry (PS100001)


Fully Sequenced ORF: >SC329812 representing NM\_001206486.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

TGGGTCCAGAAGACCTGCATATGGGCTGTTGACTCATGCAAATGA

Restriction Sites: Sgfl-Mlul

Plasmid Map:



**ACCN:** NM 001206486

Insert Size: 321 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## ATF3 (NM\_001206486) Human Untagged Clone - SC329812

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeg:** NM 001206486.2

RefSeq Size: 2010 bp RefSeq ORF: 321 bp Locus ID: 467

 UniProt ID:
 P18847

 Cytogenetics:
 1q32.3

**Protein Families:** Transcription Factors

MW: 12.3 kDa

**Gene Summary:** This gene encodes a member of the mammalian activation transcription factor/cAMP

responsive element-binding (CREB) protein family of transcription factors. This gene is induced by a variety of signals, including many of those encountered by cancer cells, and is involved in the complex process of cellular stress response. Multiple transcript variants encoding different isoforms have been found for this gene. It is possible that alternative splicing of this gene may be physiologically important in the regulation of target genes.

[provided by RefSeq, Apr 2011]

Transcript Variant: This variant (7) differs in the 5' UTR, lacks an in-frame segment of the 5' coding region, and contains an alternate terminal exon, thus differing in the 3' coding region and 3' UTR, compared to variant 1. The encoded isoform (5, also known as deltaZip2c) has a

distinct C-terminus and is shorter than isoform 1.