

Product datasheet for **SC329916**

p21 (CDKN1A) (NM_001220778) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: p21 (CDKN1A) (NM_001220778) Human Untagged Clone
Tag: Tag Free
Symbol: p21
Synonyms: CAP20; CDKN1; CIP1; MDA-6; P21; p21CIP1; SDI1; WAF1
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC329916 representing NM_001220778.
Blue=Insert sequence **Red**=Cloning site **Green**=Tag(s)

ATGTCAGAACCGGCTGGGGATGTCGTCAGAACCCATGCGGCAGCAAGGCCTGCCGCCGCTCTTCGGC
CCAGTGGACAGCGAGCAGCTGAGCCGCGACTGTGATGCGCTAATGGCGGGCTGCATCCAGGAGGCCCGT
GAGCGATGGAACCTTCGACTTTGTACCGAGACACCACTGGAGGGTGACTTCGCCTGGGAGCGTGTGCGG
GGCCTTGGCCTGCCCAAGCTCTACCTTCCACGGGGCCCCGGCGAGGCCGGGATGAGTTGGGAGGAGGC
AGGCGGCCTGGCACCTCACCTGCTCTGCTGCAGGGGACAGCAGAGGAAGACCATGTGGACCTGTCACTG
TCTTGTACCTTTGTGCCTCGCTCAGGGGAGCAGGCTGAAGGGTCCCAGGTGGACCTGGAGACTCTCAG
GGTCGAAAACGGCGGCAGACCAGCATGACAGATTCTACCACTCCAACGCCGGCTGATCTTCTCCAAG
AGGAAGCCCTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001220778
Insert Size: 495 bp
OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001220778.1
RefSeq Size:	2284 bp
RefSeq ORF:	495 bp
Locus ID:	1026
UniProt ID:	P38936
Cytogenetics:	6p21.2
Protein Families:	Druggable Genome
Protein Pathways:	Bladder cancer, Cell cycle, Chronic myeloid leukemia, ErbB signaling pathway, Glioma, Melanoma, p53 signaling pathway, Pathways in cancer, Prostate cancer
MW:	18.1 kDa
Gene Summary:	<p>This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene. [provided by RefSeq, Sep 2015]</p> <p>Transcript Variant: This variant (4) differs in the 5' UTR, lacks an in-frame portion of the 5' coding region, and initiates translation at a downstream start codon, compared to variant 3. The resulting isoform (1) has a shorter N-terminus, compared to isoform 2. Variants 1, 2, 4 and 5 encode the same isoform 1.</p>