

Product datasheet for SC330060

ZA20D3 (ZFAND6) (NM 001242912) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: ZA20D3 (ZFAND6) (NM_001242912) Human Untagged Clone

Tag: Tag Free Symbol: ZFAND6

Synonyms: AWP1; ZA20D3; ZFAND5B Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC330060 representing NM_001242912.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

ATTTGA

Restriction Sites: Sgfl-Mlul


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001242912

Insert Size: 627 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

ZA20D3 (ZFAND6) (NM_001242912) Human Untagged Clone - SC330060

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001242912.1</u>

 RefSeq Size:
 1743 bp

 RefSeq ORF:
 627 bp

 Locus ID:
 54469

 UniProt ID:
 Q6FIF0

 Cytogenetics:
 15q25.1

 MW:
 22.6 kDa

Gene Summary: Involved in regulation of TNF-alpha induced NF-kappa-B activation and apoptosis. Involved in

modulation of 'Lys-48'-linked polyubiquitination status of TRAF2 and decreases association of

TRAF2 with RIPK1. Required for PTS1 target sequence-dependent protein import into peroxisomes and PEX5 stability; may cooperate with PEX6. In vitro involved in PEX5 export

from the cytosol to peroxisomes (By similarity).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (3) differs in the 5' UTR compared to variant 1. Variants 1-7 all

encode isoform a.