

Product datasheet for SC330179

GNG2 (NM_001243773) Human Untagged Clone


Product data:

Product Type: Expression Plasmids
 Product Name: GNG2 (NM_001243773) Human Untagged Clone
 Tag: Tag Free
 Symbol: GNG2
 Vector: pCMV6-Entry (PS100001)
 Fully Sequenced ORF: >SC330179 representing NM_001243773.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
ATGGCCAGCAACAACACCGCCAGCATAGCACAAGCCAGGAAGCTGGTAGAGCAGCTTAAGATGGAAGCC
AATATCGACAGGATAAAGGTGTCCAAGGCAGCTGCAGATTTGATGGCCTACTGTGAAGCACATGCCAAG
GAAGACCCCTCCTGACCCCTGTTCCGGCTTCAGAAAACCCGTTTAGGGAGAAGAAGTTTTTCTGTGCC
ATCCTTTAA
```

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM_001243773

Insert Size: 216 bp



[View online >](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001243773.1</u>
RefSeq Size:	3899 bp
RefSeq ORF:	216 bp
Locus ID:	54331
UniProt ID:	<u>P59768</u>
Cytogenetics:	14q22.1
Protein Families:	Druggable Genome
Protein Pathways:	Chemokine signaling pathway
MW:	7.9 kDa
Gene Summary:	<p>This gene encodes one of the gamma subunits of a guanine nucleotide-binding protein. Such proteins are involved in signaling mechanisms across membranes. Various subunits forms heterodimers which then interact with the different signal molecules. [provided by RefSeq, Aug 2011]</p> <p>Transcript Variant: This variant (2) uses a different splice site in the 5' UTR, compared to variant 1. Variants 1, 2, and 3 encode the same protein. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>