

## **Product datasheet for SC330192**

## TPD52L2 (NM 001243892) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** TPD52L2 (NM\_001243892) Human Untagged Clone

Tag: Tag Free
Symbol: TPD52L2
Synonyms: D54; TPD54

Vector: pCMV6-Entry (PS100001)


Fully Sequenced ORF: >SC330192 representing NM\_001243892.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

**CCTTTCTAA** 

**Restriction Sites:** Sgfl-Mlul

Plasmid Map:





**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## TPD52L2 (NM\_001243892) Human Untagged Clone - SC330192

**ACCN:** NM 001243892

**Insert Size:** 492 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** NM 001243892.1

 RefSeq Size:
 2230 bp

 RefSeq ORF:
 492 bp

 Locus ID:
 7165

 UniProt ID:
 043399

 Cytogenetics:
 20q13.33

MW:

**Gene Summary:** This gene encodes a member of the tumor protein D52-like family. These proteins are

characterized by an N-terminal coiled-coil motif that is used to form homo- and heteromeric complexes with other tumor protein D52-like proteins. Expression of this gene may be a marker for breast cancer and acute lymphoblastic leukemia. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 12. [provided by RefSeq, Aug 2011] Transcript Variant: This variant (8) uses an alternate splice site and lacks three exons in the coding region, but maintains the reading frame, compared to variant 1. The encoded isoform (h) is shorter than isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript

alignments.

17.5 kDa