

## **Product datasheet for SC330206**

## SPIB (NM\_001243999) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** SPIB (NM\_001243999) Human Untagged Clone

Tag: Tag Free
Symbol: SPIB
Synonyms: SPI-B


**Vector:** pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC330206 representing NM\_001243999.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

**Restriction Sites:** Sgfl-Mlul

Plasmid Map:



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## SPIB (NM\_001243999) Human Untagged Clone - SC330206

**ACCN:** NM\_001243999

**Insert Size:** 534 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** NM 001243999.1

 RefSeq Size:
 3533 bp

 RefSeq ORF:
 534 bp

 Locus ID:
 6689

 UniProt ID:
 Q01892

 Cytogenetics:
 19q13.33

**Protein Families:** Transcription Factors

**MW:** 18.6 kDa

**Gene Summary:** The protein encoded by this gene is a transcriptional activator that binds to the PU-box (5'-

GAGGAA-3') and acts as a lymphoid-specific enhancer. Four transcript variants encoding

different isoforms have been found for this gene. [provided by RefSeq, Sep 2011]

Transcript Variant: This variant (3, also known as DeltaSpi-B) uses an alternate splice junction at the 3' end of a coding exon compared to variant 1, that causes a frameshift. The resulting isoform (3) has a shorter and distinct C-terminus compared to isoform 1. This isoform is found in the cytoplasm instead of the nucleus. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record

were based on transcript alignments.