

Product datasheet for SC330246

RAP1B (NM_001251917) Human Untagged Clone


Product data:

Product Type:	Expression Plasmids
Product Name:	RAP1B (NM_001251917) Human Untagged Clone
Tag:	Tag Free
Symbol:	RAP1B
Synonyms:	K-REV; RAL1B
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC330246 representing NM_001251917. Blue=Insert sequence Red=Cloning site Green=Tag(s)

ATGCGTGAGTATAAGCTAGTCGTTCTTGGCTCAGGAGGCGTTGGAAAGTCTGCTTTGGAGCAATTTACA
GCAATGAGGGATTATACATGAAAAATGGACAAGGATTTGCATTAGTTTATTCCATCACAGCACAGTCC
ACATTTAACGATTTACAAGACCTGAGAGAACAGATTCTTCGAGTTAAAGACACTGATGATGTTCCAATG
ATTCTTGTGGTAATAAGTGTGACTTGAAGATGAAAGAGTTGTAGGGAAGGAACAAGGTCAAAATCTA
GCAAGACAATGGAACAACGTGCATTCTAGAATCTTCTGCAAAATCAAAAATAAATGTTAATGAGATC
TTTTATGACCTAGTGCGGCAAATTAACAGAAAACTCCAGTGCCTGGGAAGGCTCGCAAAAAGTCATCA
TGTCAGCTGCTTAA

Restriction Sites: SgfI-MluI

Plasmid Map:



[View online »](#)

ACCN:	NM_001251917
Insert Size:	429 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001251917.1
RefSeq Size:	2046 bp
RefSeq ORF:	429 bp
Locus ID:	5908
UniProt ID:	P61224
Cytogenetics:	12q15
Protein Families:	Druggable Genome
Protein Pathways:	Chemokine signaling pathway, Focal adhesion, Leukocyte transendothelial migration, Long-term potentiation, MAPK signaling pathway, Neurotrophin signaling pathway, Renal cell carcinoma
MW:	16 kDa
Gene Summary:	<p>This gene encodes a member of the RAS-like small GTP-binding protein superfamily. Members of this family regulate multiple cellular processes including cell adhesion and growth and differentiation. This protein localizes to cellular membranes and has been shown to regulate integrin-mediated cell signaling. This protein also plays a role in regulating outside-in signaling in platelets. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 3, 5, 6 and 9. [provided by RefSeq, Oct 2011]</p> <p>Transcript Variant: This variant (3) lacks two in-frame exons in the coding region, compared to variant 1. This results in a shorter protein (isoform 2), compared to isoform 1. Variants 3 and 4 encode the same isoform (2).</p>