

# **Product datasheet for SC330252**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DSCR1L1 (RCAN2) (NM\_001251974) Human Untagged Clone

### **Product data:**

**Product Type:** Expression Plasmids

Product Name: DSCR1L1 (RCAN2) (NM\_001251974) Human Untagged Clone

Tag: Tag Free
Symbol: DSCR1L1

Synonyms: CSP2; DSCR1L1; MCIP2; RCN2; ZAKI-4; ZAKI4

**Vector:** pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC330252 representing NM\_001251974.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM 001251974

**Insert Size:** 732 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





#### **Reconstitution Method:**

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** <u>NM 001251974.1</u>

 RefSeq Size:
 3316 bp

 RefSeq ORF:
 732 bp

 Locus ID:
 10231

 UniProt ID:
 Q14206

 Cytogenetics:
 6p12.3

 MW:
 27.3 kDa

**Gene Summary:** This gene encodes a member of the regulator of calcineurin (RCAN) protein family. These

proteins play a role in many physiological processes by binding to the catalytic domain of calcineurin A, inhibiting calcineurin-mediated nuclear translocation of the transcription factor NFATC1. Expression of this gene in skin fibroblasts is upregulated by thyroid hormone, and the encoded protein may also play a role in endothelial cell function and angiogenesis. Alternatively spliced transcript variants encoding multiple isoforms have been observed for

this gene. [provided by RefSeq, Oct 2011]

Transcript Variant: This variant (2, also known as beta-1) differs in the 5' UTR and uses an alternate start codon, compared to variant 1. Variants 2 and 3 encode the same isoform (2, also known as beta), which is longer and has a distinct N-terminus, compared to isoform 1.