

Product datasheet for SC330400

UBE2L3 (NM_001256356) Human Untagged Clone


Product data:

Product Type:	Expression Plasmids
Product Name:	UBE2L3 (NM_001256356) Human Untagged Clone
Tag:	Tag Free
Symbol:	UBE2L3
Synonyms:	E2-F1; L-UBC; UBCH7; UbcM4
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC330400 representing NM_001256356. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
ATGGCGGCCAGCAGGAGGCTGATGAAGGACAACCTCCATATGATAAGGGAGCCTTCAGAATCGAAATC
AACTTTCCAGCAGAGTACCCATTCAAACCACCGAAGATCACATTTAAACAAAGATCTATCACCCAAAC
ATCGACGAAAAGGGCAGGTCTGTCTGCCAGTAATTAGTGCCGAAAACCTGGAAGCCAGCAACCAAAACC
GACCAAGTAATCCAGTCCCTCATAGCACTGGTGAATGACCCCGAGCCTGAGCACCCGCTTCGGGCTGAC
CTAGCTGAAGAATACTCTAAGGACCGTAAAAAATCTGTAAGAATGCTGAAGAGTTACAAGAAATAT
GGGAAAAGCGACCTGTGGACTAA
```

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM_001256356



[View online >](#)

Insert Size:	369 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001256356.1
RefSeq Size:	2932 bp
RefSeq ORF:	369 bp
Locus ID:	7332
UniProt ID:	P68036
Cytogenetics:	22q11.21
Protein Pathways:	Parkinson's disease, Ubiquitin mediated proteolysis
MW:	14.1 kDa
Gene Summary:	<p>The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes (E1s), ubiquitin-conjugating enzymes (E2s) and ubiquitin-protein ligases (E3s). This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is demonstrated to participate in the ubiquitination of p53, c-Fos, and the NF-kB precursor p105 in vitro. Several alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Sep 2009]</p> <p>Transcript Variant: This variant (3) uses an alternate 5' splice site, lacks an alternate exon, and initiates translation from a downstream start site, compared to variant 4. It encodes a shorter and distinct N-terminus in isoform 3, compared to isoform 4.</p>