

Product datasheet for SC330402

CCR9 (NM_001256369) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: CCR9 (NM_001256369) Human Untagged Clone

Tag: Tag Free Symbol: CCR9

Synonyms: CC-CKR-9; CDw199; GPR-9-6; GPR28

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC330402 representing NM_001256369.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

ATGTTGCTGGAGACAACCTCAGGAGCACTCTCCCTCTGA

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001256369

Insert Size: 1074 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001256369.1</u>

RefSeq Size: 2653 bp



CCR9 (NM_001256369) Human Untagged Clone - SC330402

 RefSeq ORF:
 1074 bp

 Locus ID:
 10803

 UniProt ID:
 P51686

 Cytogenetics:
 3p21.31

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction

MW: 40.7 kDa

Gene Summary: The protein encoded by this gene is a G protein-coupled receptor with seven transmembrane

domains that belongs to the beta chemokine receptor family. Chemokines and their receptors are key regulators of thymocyte migration and maturation in normal and inflammation conditions. This gene is differentially expressed in T lymphocytes of the small intestine and colon, and its interaction with chemokine 25 contributes to intestinal intraepithelial lymphocyte homing to the small intestine. This suggests a role for this gene in directing immune responses to different segments of the gastrointestinal tract. This gene and its exclusive ligand, chemokine 25, are overexpressed in a variety of malignant tumors and are closely associated with tumor proliferation, apoptosis, invasion, migration and drug

variants encoding different isoforms have been found for this gene. [provided by RefSeq, Aug 2020]

resistance. This gene maps to the chemokine receptor gene cluster. Multiple transcript

Transcript Variant: This variant (C) includes an alternate internal exon and initiates translation at a downstream, in-frame start codon, compared to variant 1. Variants B and C encode the same isoform (B), which has a shorter N-terminus compared to isoform A.