

Product datasheet for SC330446

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PBR (TSPO) (NM_001256530) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: PBR (TSPO) (NM_001256530) Human Untagged Clone

Tag: Tag Free

Symbol: PBR

Synonyms: BPBS; BZRP; DBI; IBP; MBR; mDRC; PBR; PBS; pk18; PKBS; PTBR

Mammalian Cell

Selection:

None

Vector: pCMV6-XL5

E. coli Selection: Ampicillin (100 ug/mL)

Fully Sequenced ORF: >OriGene ORF within SC111597 sequence for NM_000714 edited (data generated by NextGen

Sequencing)

TGGCGTGGGGACGGCGGCTGCCAGAGTGA

Clone variation with respect to NM_000714.4

439 a=>g

Restriction Sites: Sgfl-Mlul

ACCN: NM 001256530

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001256530.1</u>, <u>NP 001243459.1</u>

 RefSeq Size:
 1116 bp

 RefSeq ORF:
 510 bp

 Locus ID:
 706

 UniProt ID:
 P30536

 Cytogenetics:
 22q13.2

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Neuroactive ligand-receptor interaction

Gene Summary: Present mainly in the mitochondrial compartment of peripheral tissues, the protein encoded

by this gene interacts with some benzodiazepines and has different affinities than its endogenous counterpart. The protein is a key factor in the flow of cholesterol into mitochondria to permit the initiation of steroid hormone synthesis. Alternatively spliced transcript variants have been reported; one of the variants lacks an internal exon and is considered non-coding, and the other variants encode the same protein. [provided by RefSeq,

Feb 2012]

Transcript Variant: This variant (3) differs in the 5' UTR, compared to variant PBR. Variants

PBR, 3 and 4 encode the same protein.