

Product datasheet for **SC330502**

Apc10 (ANAPC10) (NM_001256706) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Apc10 (ANAPC10) (NM_001256706) Human Untagged Clone
Tag: Tag Free
Symbol: Apc10
Synonyms: APC10; DOC1
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC330502 representing NM_001256706.
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
ATGACTACACCAAACAAGACACCTCTGGTGCTGACCCCAAGCAGTTGGAAAGGACTGGAACAGTACGG
GAAATTGGGTCACAAGCTGTTTGGTCACCTCATCTTGCAAACCAGGATTTGGAGTGGATCAGTTACGA
GATGACAACTAGAAACTTATTGGCAATCAGATGGTCCAGCCTCATTTAGTGAACATCCAATTCAGA
AGAAAAACAACAGTGAAGACATTATGTATTTATGCAGACTACAAATCTGTGAAAGCTATACTCCAAGC
AAGATCTCAGTCAGAGTAGGAAATAATTTTCAACAACCTCAAGAAATTCGGCAACTTGAGTTGGTGGAA
CCAAGTGGCTGGATTCATGTTCCCTTAAGTACAATCATAAGAAGCCAACCTCGTACATTCATGATACAG
ATTGCTGTTCTAGCCAATCACCAGAATGGAAGAGACCCCATATGAGACAAATTTAAATATACACACCA
GTAGAAGAGAGCTCCATTGGTAAATTTCTAGATGTACAACCTATAGATTTTCATGATGATCGTTCAATA
AGGTGA
```

Restriction Sites: Sgfl-Mlul
ACCN: NM_001256706
Insert Size: 558 bp
OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001256706.1
RefSeq Size:	1468 bp
RefSeq ORF:	558 bp
Locus ID:	10393
UniProt ID:	Q9UM13
Cytogenetics:	4q31.21
Protein Families:	Druggable Genome
Protein Pathways:	Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation, Ubiquitin mediated proteolysis
MW:	21.3 kDa
Gene Summary:	<p>ANAPC10 is a core subunit of the anaphase-promoting complex (APC), or cyclosome, a ubiquitin protein ligase that is essential for progression through the cell cycle. APC initiates sister chromatid separation by ubiquitinating the anaphase inhibitor securin (PTTG1; MIM 604147) and triggers exit from mitosis by ubiquitinating cyclin B (CCNB1; MIM 123836), the activating subunit of cyclin-dependent kinase-1 (CDK1; MIM 116940) (summary by Wendt et al., 2001 [PubMed 11524682]).[supplied by OMIM, Feb 2011]</p> <p>Transcript Variant: This variant (1) represents the predominant transcript. Variants 1, 2, 3 and 4 encode the same isoform (1). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>