

## Product datasheet for **SC330505**

### Apc10 (ANAPC10) (NM\_001256709) Human Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Apc10 (ANAPC10) (NM\_001256709) Human Untagged Clone  
**Tag:** Tag Free  
**Symbol:** ANAPC10  
**Synonyms:** APC10; DOC1  
**Vector:** pCMV6-Entry (PS100001)  
**Fully Sequenced ORF:** >SC330505 representing NM\_001256709.  
Blue=Insert sequence Red=Cloning site Green=Tag(s)


```
ATGGGAGGTGCAAGGAAACGAATATTTTAAAAATGACTACACCAACAAGACACCTCCTGGTGCTGAC
CCCAAGCAGTTGGAAAGGACTGGAACAGTACGGGAAATTGGGTCACAAGCTGTTTGGTCACTCTCATCT
TGCAAACCAGGATTTGGAGTGGATCAGTTACGAGATGACAATCTAGAACTTATTGGCAATCAGATGGT
TCCCAGCCTCATTTAGTGAACATCCAATTCAGAAGAAAAACAACAGTGAAGACATTATGTATTTATGCA
GACTACAAATCTGATGAAAGCTATACTCCAAGCAAGATCTCAGTCAGAGTAGGAAATAATTTTCACAAC
CTTCAAGAAATTCGGCAACTTGAGTTGGTGAACCAAGTGGCTGGATTTCATGTTCCCTTAACTGACAAT
CATAAGAAGCCAACCTCGTACATTCATGATACAGATTGCTGTTCTAGCCAATCACCAGAATGGAAGAGAC
ACCCATATGAGACAAATTTAAATATACACACCAGTAGAAGAGAGCTCCATTGGTAAATTTCTAGATGT
ACAACTATAGATTTTCATGATGTATCGTTCAATAAGGTGA
```

**Restriction Sites:** Sgfl-Mlul



[View online »](#)

**Plasmid Map:**



**ACCN:** NM\_001256709

**Insert Size:** 591 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001256709.1</a>
<b>RefSeq Size:</b>	1609 bp
<b>RefSeq ORF:</b>	591 bp
<b>Locus ID:</b>	10393
<b>UniProt ID:</b>	<a href="#">Q9UM13</a>
<b>Cytogenetics:</b>	4q31.21
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation, Ubiquitin mediated proteolysis
<b>MW:</b>	22.4 kDa
<b>Gene Summary:</b>	<p>ANAPC10 is a core subunit of the anaphase-promoting complex (APC), or cyclosome, a ubiquitin protein ligase that is essential for progression through the cell cycle. APC initiates sister chromatid separation by ubiquitinating the anaphase inhibitor securin (PTTG1; MIM 604147) and triggers exit from mitosis by ubiquitinating cyclin B (CCNB1; MIM 123836), the activating subunit of cyclin-dependent kinase-1 (CDK1; MIM 116940) (summary by Wendt et al., 2001 [PubMed 11524682]).[supplied by OMIM, Feb 2011]</p> <p>Transcript Variant: This variant (5) differs in the 5' UTR and initiates translation at an alternate start codon, compared to variant 1. The encoded isoform (2) is longer and has a distinct N-terminus, compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>