

## **Product datasheet for SC330616**

## CD63 (NM\_001257400) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** CD63 (NM\_001257400) Human Untagged Clone

Tag: Tag Free Symbol: CD63

Synonyms: LAMP-3; ME491; MLA1; OMA81H; TSPAN30

**Vector:** pCMV6-Entry (PS100001)


Fully Sequenced ORF: >SC330616 representing NM\_001257400.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

ATTGTCTTTGCCTGCTCGTGAAGAGTATCAGAAGTGGCTACGAGGTGATGTAG

**Restriction Sites:** Sgfl-Mlul

Plasmid Map:





**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## CD63 (NM\_001257400) Human Untagged Clone - SC330616

**ACCN:** NM\_001257400

**Insert Size:** 471 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 001257400.1</u>

 RefSeq Size:
 806 bp

 RefSeq ORF:
 471 bp

 Locus ID:
 967

 UniProt ID:
 P08962

 Cytogenetics:
 12q13.2

**Protein Families:** Druggable Genome, Transmembrane

Protein Pathways: Lysosome MW: 17.3 kDa

**Gene Summary:** The protein encoded by this gene is a member of the transmembrane 4 superfamily, also

known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different

protein isoforms. [provided by RefSeq, Apr 2012]

Transcript Variant: This variant (8) contains a distinct 5' UTR exon and lacks an in-frame portion of the 5' coding region, compared to variant 1. The resulting isoform (D) has a shorter

N-terminus, compared to isoform A. Variants 8 and 9 encode the same isoform D.