

Product datasheet for **SC330691**

RPS3 (NM_001260506) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	RPS3 (NM_001260506) Human Untagged Clone
Tag:	Tag Free
Symbol:	RPS3
Synonyms:	S3
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC330691 representing NM_001260506. Blue=Insert sequence Red=Cloning site Green=Tag(s)

ATGGCAGTGCAAATATCCAAGAAGAGGAAGTTTGTGCTGATGGCATCTTCAAAGCTGAAGTGAATGAG
 TTTCTTACTCGGGAGCTGGCTGAAGATGGCTACTCTGGAGTTGAGGTGCGAGTTACCAACCAGGACA
 GAAATCATTATCTTAGCCACCAGAACACAGAATGTTCTTGGTGAGAAGGGCCGGCGGATTCGGGAACGTG
 ACTGCTGTAGTTCAGAAGAGGTTTGGCTTTCCAGAGGGCAGTGTAGAGTTGAAAATAATGGTGATGGTA
 ACAGGATATCCCTTGCTTCCTTTAAAGCTTTATGCTGAAAAGGTGGCCACTAGAGGTCTGTGTGCCATT
 GCCCAGGCAGAGTCTCTGCGTTACAACTCCTAGGAGGGCTTGCTGTGCGGAGGGCCTGCTATGGTGTG
 CTGCGGTTTCATCATGGAGAGTGGGGCCAAAGGCTGCGAGGTTGTGGTGTCTGGGAACTCCGAGGACAG
 AGGGCTAAATCCATGAAGTTTGTGGATGGCCTGATGATCCACAGCGGAGACCCTGTAACTACTACGTT
 GACACTGCTGTGCGCCACGTGTTGCTCAGACAGGGTGTGCTGGGCATCAAGGTGAAGATCATGCTGCCC
 TGGGACCAACTGGTAAGATTGGCCCTAAGAAGCCCCTGCCTGACCACGTGAGCATTGTGGAACCCAAA
 GATGAGATACTGCCCACCACCCCATCTCAGAACAGAAGGTTGGGAAGCCAGAGCCGCTGCCATGCCC
 CAGCCAGTCCCCACAGCA**TAA**

Restriction Sites:	Sgfl-MluI
ACCN:	NM_001260506
Insert Size:	780 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).


[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001260506.1](#)

RefSeq Size: 2187 bp

RefSeq ORF: 780 bp

Locus ID: 6188

UniProt ID: [P23396](#)

Cytogenetics: 11q13.4

Protein Pathways: Ribosome

MW: 28.5 kDa

Gene Summary: Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 40S subunit, where it forms part of the domain where translation is initiated. The protein belongs to the S3P family of ribosomal proteins. Studies of the mouse and rat proteins have demonstrated that the protein has an extraribosomal role as an endonuclease involved in the repair of UV-induced DNA damage. The protein appears to be located in both the cytoplasm and nucleus but not in the nucleolus. Higher levels of expression of this gene in colon adenocarcinomas and adenomatous polyps compared to adjacent normal colonic mucosa have been observed. This gene is co-transcribed with the small nucleolar RNA genes U15A and U15B, which are located in its first and fifth introns, respectively. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2012]

Transcript Variant: This variant (3) has two additional segments, one in the coding region and one in the 3' UTR, compared to variant 1. The resulting isoform (2) is longer and has an additional segment in the central region, compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.