

Product datasheet for **SC330713**

TXNRD1 (NM_001261445) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: TXNRD1 (NM_001261445) Human Untagged Clone
Tag: Tag Free
Symbol: TXNRD1
Synonyms: GRIM-12; TR; TR1; TRXR1; TXNR
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC330713 representing NM_001261445.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```


ATGGAGGACGGTCGGGCCCTGGAAGGAACGCTCTCGGAATTGGCCGCGGAAACCGATCTGCCCGTTGTG
TTTGTGAAACAGAGAAAGATAGGCGGCCATGGTCCAACCTTGAAGGCTTATCAGGAGGGCAGACTTCAA
AAGCTACTAAAAATGAACGGCCCTGAAGATCTTCCAAGTCTATGACTATGACCTTATCATCATTGGA
GGTGGCTCAGGAGGTCTGGCAGTGCTAAGGAGGCAGCCCAATATGGCAAGAAGGTGATGGTCTGGAC
TTTGTCACTCCCACCCCTCTTGGAACTAGATGGGGTCTCGGAGGAACATGTGTGAATGTGGGTTGCATA
CCTAAAAAAGTATGATCAAGCAGCTTGTAGGACAAGCCCTGCAAGACTCTCGAAATATGGATGG
AAAGTCGAGGAGACAGTTAAGCATGATTGGGACAGAATGATAGAAGCTGTACAGAATCACATTGGCTCT
TTGAATTGGGGCTACCGAGTAGCTCTGCGGGAGAAAAAGTCGTCTATGAGAATGCTTATGGGCAATTT
ATTGGTCTCACAGGATTAAGGCAACAAAATAATAAAGGCAAGAAAAATTTATTCAGCAGAGAGATTT
CTCATTGCCACTGGTGAAGACCACGTTACTTGGGCATCCCTGGTGACAAAGAATACTGCATCAGCAGT
GATGATCTTTTCTCCTTGCCTTACTGCCCAGGTAAGACCTGGTGTGGAGCATCCTATGTCGCTTTG
GAGTGCCTGGATTTCTTGTCTGTTTGGTTTAGACGTCCTGTTATGGTTAGGTCCATTCTTCTTAGA
GGATTTGACCAGGACATGGCCAACAAAATTGGTGAACACATGGAAGAACATGGCATCAAGTTTATAAGA
CAGTTCGTACCAATTAAGTTGAACAAAATTGAAGCAGGGACACCCAGGCCGACTCAGAGTAGTAGCTCAG
TCCACCAATAGTGAGGAAATCATTGAAGGAGAATAAATACGGTGATGCTGGCAATAGGAAGAGATGCT
TGCAACAAGAAAAATTGGCTTAGAAAACCGTAGGGGTGAAGATAAATGAAAAGACTGGAAAAATACCTGTC
ACAGATGAAGAACAGACCAATGTGCCTTACATCTATGCCATTGGCGATATATTGGAGGATAAGGTGGAG
CTCACCCAGTTGCAATCCAGGCAGGAAGATTGCTGGCTCAGAGGCTCTATGCAGTTTCCACTGTCAAG
TGTGACTATGAAAATGTTCCAACCTGTATTTACTCCTTTGGAATATGGTGCTTGTGGCCTTTCTGAG
GAGAAAGCTGTGGAGAAGTTTGGGGAAGAAAATTTGAGGTTTACCATAGTTACTTTTGGCCATTGGAA
TGGACGATCCGTC AAGAGATAACAACAAATGTTATGCAAAAATAATCTGTAATACTAAAGACAATGAA
CGTGTGTGGGCTTTACGTAAGTGGTCCAATGCTGGAGAAGTTACACAAGGCTTTGCAGCTGCGCTC
AAATGTGGACTGACCAAAAAGCAGCTGGACAGCAAAATGGAATCCACCTGTCTGTGCAGAGGTATTC
ACAACATTGTCTGTGACCAAGCGCTCTGGGGCAAGCATCTCCAGGCTGGCTGCTGAGGTTAA
  
```

Restriction Sites: SgfI-MluI



[View online »](#)

Plasmid Map:



ACCN: NM_001261445

Insert Size: 1650 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001261445.1](#)

RefSeq Size: 4291 bp

RefSeq ORF: 1650 bp

Locus ID: 7296

UniProt ID: [Q16881](#)

Cytogenetics: 12q23.3

Protein Families: Druggable Genome

Protein Pathways: Pyrimidine metabolism

MW: 60.2 kDa

Gene Summary: The protein encoded by this gene belongs to the pyridine nucleotide-disulfide oxidoreductase family, and is a member of the thioredoxin (Trx) system. Three thioredoxin reductase (TrxR) isozymes are found in mammals. TrxRs are selenocysteine-containing flavoenzymes, which reduce thioredoxins, as well as other substrates, and play a key role in redox homeostasis. This gene encodes an ubiquitously expressed, cytosolic form of TrxR, which functions as a homodimer containing FAD, and selenocysteine (Sec) at the active site. Sec is encoded by UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, the Sec insertion sequence (SECIS) element, which is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. Alternative splicing, primarily at the 5' end, results in transcript variants encoding same or different isoforms, including a glutaredoxin-containing isoform that is predominantly expressed in testis. [provided by RefSeq, May 2017]

Transcript Variant: This variant (5) uses an alternate donor splice site at the 5' terminal exon, which results in translation initiation from an in-frame upstream start codon compared to variant 1. The encoded isoform (3) has a longer and distinct N-terminus compared to isoform 1.