

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC330739

MRG15 (MORF4L1) (NM_001265603) Human Untagged Clone

Product data:

RF4L1) (NM_001265603) Human Untagged Clone
6; HsT17725; MEAF3; MORFRG15; MRG15; S863-6
/ (PS100001)
epresenting NM_001265603. sequence Red=Cloning site Green=Tag(s) GCCCCAGGAAAGAAGACATCTGGTCTGCAACAGAAAAATGTTGAAGTGAAAACGAAA GAAAACACCTGGAAATGGAGATGGTGGCAGTACCAGTGAGACCCCTCAGCCTCCTCGG CCGGGTAGATCCTACTGTTGAAAATGAGGAAACATTCATGAACAGAGTTGAAGTTAAA GAAGAGCTAAAACCGTGGCTTGTTGATGACTGGGACTTAATTACCAGGCAAAAACAG TCCTGCCAAGAAGAATGTGGATTCCATTCTTGAGGGATTATGCAAATTACAAGAAATCT AGATAATAAGGAGTATGCGGTTAATGAAGGTGTGGGCAGGGATAAAAGAAATACTTCAAC TACCCAGCTACTCTATAAATTTGAGAGACCACAGTATGCTGAAATTCTTGCAGATCAT CATGTCCCAGGTGATGGGAGCGCCACATCTCCTGAGATTATTTGTACGAAATTCTGCAGATCAT CATGTCCCAGGTGTATGGAGCGCCCACATCTCCTGAGATTATTTGTACGAAATTGGAGCA TACCCCGGTGATGGAGAGAGCCTTGCTTTATTACTCAATTACTTCACGATTGCAA

Restriction Sites:


Sgfl-Mlul



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Plasmid Map:



ACCN: Insert Size:	NM_001265603 708 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

SC330739 ORIGENE MRG15 (MORF4L1) (NM_001265603) Human Untagged Clone – SC330739

Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001265603.1</u>
RefSeq Size:	1779 bp
RefSeq ORF:	708 bp
Locus ID:	10933
UniProt ID:	<u>Q9UBU8</u>
Cytogenetics:	15q25.1
Protein Families:	Transcription Factors
MW:	26.7 kDa
Gene Summary:	Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the mSin3A complex which acts to repress transcription by deacetylation of nucleosomal histones.

Required for homologous recombination repair (HRR) and resistance to mitomycin C (MMC). Involved in the localization of PALB2, BRCA2 and RAD51, but not BRCA1, to DNA-damage foci. [UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (3) lacks two consecutive internal exons and initiates translation at a downstream, in-frame start codon, compared to variant 1. Variants 3, 4 and 5 encode the same isoform (3), which has a shorter N-terminus compared to isoform 1.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US