

# **Product datasheet for SC330801**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HSD11B1L (NM\_001267871) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** HSD11B1L (NM\_001267871) Human Untagged Clone

Tag: Tag Free
Symbol: HSD11B1L

Synonyms: 11-beta-HSD3; 11-DH3; HSD1L; HSD3; SCDR10; SCDR10B; SDR26C2

**Vector:** pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC330801 representing NM\_001267871.


Blue=Insert sequence Red=Cloning site Green=Tag(s)

TTTGGCCATGATTGA

**Restriction Sites:** Sgfl-Mlul



#### Plasmid Map:



**ACCN:** NM\_001267871

**Insert Size:** 705 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** NM 001267871.1

 RefSeq Size:
 1554 bp

 RefSeq ORF:
 705 bp

 Locus ID:
 374875

 UniProt ID:
 Q7Z5J1

 Cytogenetics:
 19p13.3

**Protein Families:** Druggable Genome

MW: 25.8 kDa

**Gene Summary:** This gene is a member of the hydroxysteroid dehydrogenase family. The encoded protein is

similar to an enzyme that catalyzes the interconversion of inactive to active glucocorticoids (e.g. cortisone). Alternatively spliced transcript variants encoding multiple isoforms have been

observed for this gene. [provided by RefSeq, Jun 2012]

Transcript Variant: This variant (i) has multiple differences in the coding region, one of which results in initiation of translation at a downstream in-frame start codon, compared to variant g. An additional difference results in a frameshift. The encoded protein (isoform i) has a

distinct C-terminus and is shorter than isoform g.