

Product datasheet for SC330803

BORIS (CTCFL) (NM_001269050) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	BORIS (CTCFL) (NM_001269050) Human Untagged Clone
Tag:	Tag Free
Symbol:	BORIS
Synonyms:	BORIS; CT27; CTCF-T; dJ579F20.2; HMGB1L1
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC330803 representing NM_001269050. Blue=Insert sequence Red=Cloning site Green=Tag(s)
	ATGTCAGGAGATGAAAGAAGTGACGAAATTGTTCTCACAGTTTCAAATTCAAATGTGGAAGAACAAGAG GATCAACCTACAGCTGGTCAAGCAGATGCTGAAAAGGCCAAATCTACAAAAAATCAAAGAAAG
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001269050
Insert Size:	1269 bp



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GRIGENE BORIS (CTCFL) (NM_001269050) Human Untagged Clone – SC330803	
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	 Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM 001269050.1</u>
RefSeq Size:	2302 bp
RefSeq ORF:	1269 bp
Locus ID:	140690
UniProt ID:	<u>Q8NI51</u>
Cytogenetics:	20q13.31
Protein Families:	Transcription Factors
MW:	48.1 kDa
Gene Summary:	CCCTC-binding factor (CTCF), an 11-zinc-finger factor involved in gene regulation, utilizes different zinc fingers to bind varying DNA target sites. CTCF forms methylation-sensitive insulators that regulate X-chromosome inactivation. This gene is a paralog of CTCF and appears to be expressed primarily in the cytoplasm of spermatocytes, unlike CTCF which is expressed primarily in the nucleus of somatic cells. CTCF and the protein encoded by this gene are normally expressed in a mutually exclusive pattern that correlates with resetting of methylation marks during male germ cell differentiation. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2012] Transcript Variant: This variant (12, also known as B4) differs in the 5' and 3' UTRs and has

Transcript Variant: This variant (12, also known as B4) differs in the 5' and 3' UTRs and has multiple coding region differences, compared to variant 1. The translation initiation is at a downstream in-frame AUG and the resulting isoform (10, also known as B4) has a shorter N-terminus and a shorter and distinct C-terminus, compared to isoform 1.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US