

Product datasheet for **SC330834**

CDK2AP1 (NM_001270434) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: CDK2AP1 (NM_001270434) Human Untagged Clone
Tag: Tag Free
Symbol: CDK2AP1
Synonyms: doc-1; DOC1; DORC1; p12DOC-1; ST19
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC330834 representing NM_001270434.
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
ATGGCAACGTCTTCACAGTACCGCCAGCTGCTCAGTGACTACGGGCCACCGTCCCTAGGCTACACCCAG
GGAAGTGGGAACAGCCAGGTGCCCAAAGCAAATACGCGGAGCTGCTGGCCATCATTGAAGAGCTGGGG
AAGGAGATCAGACCCACGTACGCAGGGAGCAAGAGTGCCATGGAGAGGCTGAAGCGCGGCATCATTAC
GCTAGAGGACTGGTTCGGGAGTGCTTGGCAGAAACGGAACGGAATGCCAGATCCTAG
```

Restriction Sites: SgfI-MluI
ACCN: NM_001270434
Insert Size: 264 bp
OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001270434.1](#)



[View online »](#)

RefSeq Size: 1164 bp

RefSeq ORF: 264 bp

Locus ID: 8099

UniProt ID: [O14519](#)

Cytogenetics: 12q24.31

MW: 9.6 kDa

Gene Summary: The protein encoded by this gene is a cyclin-dependent kinase 2 (CDK2) -associated protein which is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. This protein was found to also interact with DNA polymerase alpha/primase and mediate the phosphorylation of the large p180 subunit, which suggests a regulatory role in DNA replication during the S-phase of the cell cycle. This protein also forms a core subunit of the nucleosome remodeling and histone deacetylation (NURD) complex that epigenetically regulates embryonic stem cell differentiation. This gene thus plays a role in both cell-cycle and epigenetic regulation. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2012]

Transcript Variant: This variant (3) differs in the 5' UTR and coding region and represents the use of an alternate promoter, compared to variant 1. This difference results in the use of an in-frame downstream start codon and a protein (isoform 2) with a shorter N-terminus, compared to isoform 1. Variants 2 and 3 encode the same protein (isoform 2).