

Product datasheet for SC330896

KLF7 (NM 001270943) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: KLF7 (NM_001270943) Human Untagged Clone

Tag: Tag Free
Symbol: KLF7
Synonyms: UKLF

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC330896 representing NM_001270943.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

TGTTTTTCCAGGTCTGACCATCTTGCCCTCCACATGAAGAGACATATCTAA

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001270943

Insert Size: 810 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 001270943.1

 RefSeq Size:
 7990 bp

 RefSeq ORF:
 810 bp

 Locus ID:
 8609

 UniProt ID:
 075840

 Cytogenetics:
 2q33.3

Protein Families: Transcription Factors

MW: 29.6 kDa

Gene Summary: The protein encoded by this gene is a member of the Kruppel-like transcriptional regulator

contain three C2H2 zinc fingers at the C-terminus that mediate binding to GC-rich sites. This protein may contribute to the progression of type 2 diabetes by inhibiting insulin expression and secretion in pancreatic beta-cells and by deregulating adipocytokine secretion in adipocytes. A pseudogene of this gene is located on the long arm of chromosome 3. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2012] Transcript Variant: This variant (3) differs in the 5' UTR and initiates translation at an alternate start codon, compared to variant 1. The resulting protein (isoform 3) has a distinct N-terminus and is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript

and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript

family. Members in this family regulate cell proliferation, differentiation and survival and

alignments.