

Product datasheet for SC330980

PPP2R4 (PTPA) (NM 001271832) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: PPP2R4 (PTPA) (NM_001271832) Human Untagged Clone

Tag: Tag Free Symbol: PTPA

Synonyms: PP2A; PPP2R4; PR53

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC330980 representing NM_001271832.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

Restriction Sites: Sgfl-Mlul




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001271832

Insert Size: 885 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.


3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.



RefSeq: <u>NM 001271832.1</u>

 RefSeq Size:
 2677 bp

 RefSeq ORF:
 885 bp

 Locus ID:
 5524

 UniProt ID:
 Q15257

 Cytogenetics:
 9q34.11

Protein Families: Druggable Genome, Phosphatase

MW: 33.5 kDa

Gene Summary: Protein phosphatase 2A is one of the four major Ser/Thr phosphatases and is implicated in

the negative control of cell growth and division. Protein phosphatase 2A holoenzymes are heterotrimeric proteins composed of a structural subunit A, a catalytic subunit C, and a regulatory subunit B. The regulatory subunit is encoded by a diverse set of genes that have been grouped into the B/PR55, B'/PR61, and B''/PR72 families. These different regulatory subunits confer distinct enzymatic specificities and intracellular localizations to the holozenzyme. The product of this gene belongs to the B' family. This gene encodes a specific phosphotyrosyl phosphatase activator of the dimeric form of protein phosphatase 2A. Alternative splicing results in multiple transcript variants encoding different isoforms.

[provided by RefSeq, Jul 2008]

Transcript Variant: This variant (7) lacks two consecutive exons in the coding region, compared to variant 1. The resulting isoform (f) lacks an internal segment, compared to isoform a. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.