

Product datasheet for **SC331003**

HSPC210 (GSKIP) (NM_001271906) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: HSPC210 (GSKIP) (NM_001271906) Human Untagged Clone
Tag: Tag Free
Symbol: HSPC210
Synonyms: C14orf129; HSPC210
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC331003 representing NM_001271906.
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
ATGGAACAGACTGTAATCCCATGGAGCTAAGCAGTATGTCAGGATTTGAAGAAGGTTTCAGAGCTGAAC
GGTTTTGAAGGAAGTACATGAAAGACATGAGGCTCGAAGCTGAAGCAGTTGTAATGATGTTCTCTTT
GCTGTTAAACAACATGTTTGTCTCGAAAAGCCTGCGGTGTGCGGATGATGTGCCTATATCAATGTGGAA
ACAAAGGAAAGAAACAGATATTGCCTAGAAGTCACTGAAGCAGGGCTCAAGGTGGTAGGCTATGCTTTT
GACCAGGTAGATGATCATTTACAGACTCCCTACCATGAAACAGTCTACTCCTTGTGGATACACTCAGC
CCCGCCTACCGAGAAGCATTTGGAAACGCACTGCTTCAAAGACTGGAAGCTTTGAAAAGAGATGGACAG
TCATGA
```

Restriction Sites: SgfI-MluI
ACCN: NM_001271906
Insert Size: 420 bp
OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001271906.1](#)

RefSeq Size: 2150 bp

RefSeq ORF: 420 bp

Locus ID: 51527

UniProt ID: [Q9P0R6](#)

Cytogenetics: 14q32.2

MW: 15.6 kDa

Gene Summary: This gene encodes a protein that is involved as a negative regulator of GSK3-beta in the Wnt signaling pathway. The encoded protein may play a role in the retinoic acid signaling pathway by regulating the functional interactions between GSK3-beta, beta-catenin and cyclin D1, and it regulates the beta-catenin/N-cadherin pool. The encoded protein contains a GSK3-beta interacting domain (GID) in its C-terminus, which is similar to the GID of Axin. The protein also contains an evolutionarily conserved RII-binding domain, which facilitates binding with protein kinase-A and GSK3-beta, enabling its role as an A-kinase anchoring protein. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, Dec 2012]

Transcript Variant: This variant (4) differs in the 5' UTR compared to variant 1. Variants 1, 2, 3 and 4 encode the same protein.