

## **Product datasheet for SC331321**

## NCR2 (NM 001199509) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** NCR2 (NM\_001199509) Human Untagged Clone

Tag: Tag Free Symbol: NCR2

**Synonyms:** CD336; dJ149M18.1; LY95; NK-p44; NKP44

**Vector:** pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC331321 representing NM\_001199509.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

CACTTCCCACTGAGCCACAGGGCACCAGGGGGGACATATGGTGGAAAACCATGA

Restriction Sites: Sgfl-Mlul

**ACCN:** NM\_001199509

**Insert Size:** 813 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** NM 001199509.1

 RefSeq Size:
 1084 bp

 RefSeq ORF:
 813 bp

 Locus ID:
 9436

 UniProt ID:
 095944

 Cytogenetics:
 6p21.1

Protein Families: Druggable Genome, Transmembrane
Protein Pathways: Natural killer cell mediated cytotoxicity

MW: 29.7 kDa

**Gene Summary:** Cytotoxicity-activating receptor that may contribute to the increased efficiency of activated

natural killer (NK) cells to mediate tumor cell lysis.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) has an additional in-frame segment and an additional exon which results in frame-shift, as compared to variant 1. The resulting isoform (2) has an additional internal segment and a shorter and distinct C-terminus, as compared to isoform 1. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to

make the sequence consistent with the reference genome assembly. The genomic

coordinates used for the transcript record were based on alignments.