

Product datasheet for SC331622

CCN4 (NM 001204869) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: CCN4 (NM_001204869) Human Untagged Clone

Tag: Tag Free Symbol: CCN4

Synonyms: WISP1; WISP1-OT1; WISP1-UT1; WISP1c; WISP1tc

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC331622 representing NM_001204869.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

CATCAGCACACGCTCCTATCAACCCAAGTACTGTGGAGTTTGCATGGACAATAG

Restriction Sites: Sgfl-Mlul

ACCN: NM_001204869

Insert Size: 468 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 001204869.1

 RefSeq Size:
 4739 bp

 RefSeq ORF:
 468 bp

 Locus ID:
 8840

 UniProt ID:
 095388

 Cytogenetics:
 8q24.22

Protein Families: Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS,

Secreted Protein, Stem cell relevant signaling - DSL/Notch pathway, Stem cell relevant

signaling - Wnt Signaling pathway

MW: 16.8 kDa

Gene Summary: This gene encodes a member of the WNT1 inducible signaling pathway (WISP) protein

subfamily, which belongs to the connective tissue growth factor (CTGF) family. WNT1 is a member of a family of cysteine-rich, glycosylated signaling proteins that mediate diverse developmental processes. The CTGF family members are characterized by four conserved cysteine-rich domains: insulin-like growth factor-binding domain, von Willebrand factor type C module, thrombospondin domain and C-terminal cystine knot-like domain. This gene may be downstream in the WNT1 signaling pathway that is relevant to malignant transformation. It is expressed at a high level in fibroblast cells, and overexpressed in colon tumors. The encoded protein binds to decorin and biglycan, two members of a family of small leucine-rich proteoglycans present in the extracellular matrix of connective tissue, and possibly prevents the inhibitory activity of decorin and biglycan in tumor cell proliferation. It also attenuates p53-mediated apoptosis in response to DNA damage through activation of the Akt kinase. It is 83% identical to the mouse protein at the amino acid level. Multiple alternatively spliced transcript variants have been identified. [provided by RefSeq, Mar 2011]

Transcript Variant: This variant (3) lacks two consecutive exons in the coding region, as compared to variant 1. The resulting isoform (3) has a shorter and distinct C-terminus, as compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript

alignments.