

Product datasheet for SC331625

RAB43 (NM 001204883) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: RAB43 (NM_001204883) Human Untagged Clone

Tag: Tag Free Symbol: RAB43

Synonyms: RAB11B; RAB41

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC331625 representing NM_001204883.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

TGGGGCTGCGGGTGCTGA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001204883

Insert Size: 639 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001204883.1</u>

 RefSeq Size:
 4232 bp

 RefSeq ORF:
 639 bp

 Locus ID:
 339122

 UniProt ID:
 Q86YS6

 Cytogenetics:
 3q21.3

Protein Families: Druggable Genome

MW: 23.3 kDa

Gene Summary: The small GTPases Rab are key regulators of intracellular membrane trafficking, from the

formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. The low intrinsic GTPase activity of RAB43 is activated by USP6NL. Involved in retrograde transport from the endocytic pathway to the Golgi apparatus. Involved in the transport of Shiga toxin from early and recycling endosomes to the trans-Golgi network. Required for the structural integrity of the Golgi complex. Plays a role in the maturation of phagosomes that engulf pathogens, such as S.aureus and M.tuberculosis.

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) differs in the 5' UTR compared to variant 1. Variants 1-5 all encode the same isoform (a). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.