

Product datasheet for **SC331771**

Ikaros (IKZF1) (NM_001220766) Human Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Ikaros (IKZF1) (NM_001220766) Human Untagged Clone
Tag: Tag Free
Symbol: Ikaros
Synonyms: hlk-1; Hs.54452; IK1; IKAROS; LYF1; PRO0758; ZNFN1A1
Vector: pCMV6-Entry (PS100001)
Fully Sequenced ORF: >SC331771 representing NM_001220766.
Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
ATGGATGCTGATGAGGGTCAAGACATGTCCCAAGTTTCAGGGAAGGAAAGCCCCCTGTAAGCGATACT
CCAGATGAGGGCGATGAGCCCATGCCGATCCCCGAGGACCTCTCCACCACCTCGGGAGGACAGCAAAGC
TCCAAGAGTGACAGAGTCGTGGGAGAACGGCCCTTCCAGTGCAATCAGTGCGGGGCTCATTACCCAG
AAGGGCAACCTGCTCCGGCACATCAAGTGCATTCCGGGGAGAAGCCCTCAAATGCCACCTCTGCAAC
TACGCCTGCCCGGAGGACGCCCTCACTGGCCACCTGAGGACGCACTCCGTTGGTAAACCTCACAAA
TGTGGATATTGTGGCCGAAGCTATAAACAGCGAAGCTCTTTAGAGGAACATAAAGAGCGTGCACAAAC
TACTTGGAAAGCATGGGCCTTCCGGGCACACTGTACCCAGTCATTAAGAAGAACTAATCACAGTGAA
ATGGCAGAAGACCTGTGCAAGATAGGATCAGAGAGATCTCTCGTGTGGCAGACTAGCAAGTAACGTC
GCCAAACGTAAGAGCTCTATGCCTCAGAAATTTCTTGGGGACAAGGGCTGTCCGACACGCCCTACGAC
AGCAGCGCCAGCTACGAGAAGGAGAACGAAATGATGAAGTCCCACGTGATGGACCAAGCCATCAACAAC
GCCATCAACTACCTGGGGCCGAGTCCCTGCGCCCGCTGGTGCAGACGCCCCGGGGCGTTCCGAGGTG
GTCCCGGTCATCAGCCCGATGTACCAGCTGCACAAGCCGCTCGCGGAGGGCACCCCGCGCTCCAACCAC
TCGGCCCAGGACAGCGCCGTGGAGAACCTGCTGCTCTCCAAGGCAAGTTGGTGCCTCGGAGCGC
GAGGCGTCCCCGAGCAACAGCTGCCAAGACTCCACGGACACCCGAGAGCAACAACGAGGAGCAGCGCAGC
GGTCTCATCTACCTGACCAACCACATCGCCCCGACGCGCAACGGGCTGTCTGCTCAAGGAGGAGCAC
CGCGCCTACGACCTGCTGCGCGCCGCTCCGAGAAGTCCGAGGACGCGCTCCGCGTGGTACGACACCAGC
GGGGAGCAGATGAAGGTGTACAAGTGCGAACACTGCCGGGTGCTTCTTGGATCAGCTCATGTACACC
ATCCACATGGGCTGCCACGGCTCCGTGATCCTTTTGTGAGTGAACATGTGCGGCTACCACAGCCAGGAC
CGGTACGATTCTCGTCGCACATAACGCGAGGGGAGCACCGCTTCCACATGAGCTAA
```

Restriction Sites: SgfI-MluI
ACCN: NM_001220766
Insert Size: 1299 bp



[View online »](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001220766.1
RefSeq Size:	5956 bp
RefSeq ORF:	1299 bp
Locus ID:	10320
Cytogenetics:	7p12.2
Protein Families:	Druggable Genome, Transcription Factors
MW:	48.3 kDa
Gene Summary:	<p>This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemo-lymphopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding to a specific core DNA sequence element in the promoters of target genes. The non-DNA-binding isoforms are largely found in the cytoplasm, and are thought to function as dominant-negative factors. Overexpression of some dominant-negative isoforms have been associated with B-cell malignancies, such as acute lymphoblastic leukemia (ALL). [provided by RefSeq, May 2014]</p> <p>Transcript Variant: This variant (3) lacks an in-frame coding exon in the 5' region compared to variant 1. This results in a shorter isoform (3, also known as Ik-2 as described by Ezzat et al. in PMID:12937159) missing an internal protein segment compared to isoform 1. This isoform is localized in the nucleus, and contains requisite three N-terminal zinc finger motifs for high affinity DNA binding (PMID:9892693).</p>