

Product datasheet for SC331884

DEF8 (NM 001242821) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: DEF8 (NM_001242821) Human Untagged Clone

Tag: Tag Free Symbol: DEF8

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC331884 representing NM_001242821.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GACCAAAGTTCCTGCCTCCGCTGGGCTCACATTCAGATGTGA

Restriction Sites: Sgfl-Rsrll

ACCN: NM_001242821

Insert Size: 594 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001242821.1</u>

 RefSeq Size:
 835 bp

 RefSeq ORF:
 594 bp

 Locus ID:
 54849

 UniProt ID:
 Q6ZN54

 Cytogenetics:
 16q24.3

 MW:
 23 kDa

Gene Summary: Positively regulates lysosome peripheral distribution and ruffled border formation in

osteoclasts. Involved in bone resorption.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (8) has multiple differences in the 5' and 3' UTRs, compared to variant 1. These differences cause translation initiation at an in-frame downstream start codon, compared to variant 1. The encoded protein (isoform 2) has shorter N- and C-termini,

compared to isoform 1. Variants 2, 8, and 9 encode the same protein.