

## Product datasheet for **SC332076**

### **BID (NM\_001244572) Human Untagged Clone**

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** BID (NM\_001244572) Human Untagged Clone  
**Tag:** Tag Free  
**Symbol:** BID  
**Synonyms:** FP497  
**Vector:** pCMV6-Entry (PS100001)  
**Fully Sequenced ORF:** >SC332076 representing NM\_001244572.  
**Blue**=Insert sequence **Red**=Cloning site **Green**=Tag(s)

```
ATGGACCGTAGCATCCCTCCGGCCTGGTGAACGGCCTGGCCCTGCAGCTCAGGAACACCAGCCGGTCC
GAGGAGGACCGGAACAGGGACCTGGCCACTGCCCTGGAGCAGCTGCTGCAGGCCTACCCTAGAGACATG
GAGAAGGAGAAGACCATGCTGGTCTGGCCCTGCTGCTGGCCAAGAAGGTGGCCAGTCACACGCCGTCC
TTGCTCCGTGATGCTTTTACACAACAGTGAATTTTATAACCAGAACCTACGCACCTACGTGAGGAGC
TTAGCCAGAAATGGGATGGACTGA
```

**Restriction Sites:** Sgfl-Mlul  
**ACCN:** NM\_001244572  
**Insert Size:** 300 bp  
**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).  
**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  
**Reconstitution Method:**  

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

  
**RefSeq:** [NM\\_001244572.1](#)



[View online »](#)

RefSeq Size:	2254 bp
RefSeq ORF:	300 bp
Locus ID:	637
UniProt ID:	<a href="#">P55957</a>
Cytogenetics:	22q11.21
Protein Families:	Druggable Genome
Protein Pathways:	Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Apoptosis, Natural killer cell mediated cytotoxicity, p53 signaling pathway, Pathways in cancer, Viral myocarditis
MW:	11.3 kDa
Gene Summary:	<p>This gene encodes a death agonist that heterodimerizes with either agonist BAX or antagonist BCL2, and thus regulate apoptosis. The encoded protein is a member of the BCL-2 family of cell death regulators. It is a mediator of mitochondrial damage induced by caspase-8 (CASP8); CASP8 cleaves this encoded protein, and the COOH-terminal part translocates to mitochondria where it triggers cytochrome c release. Multiple alternatively spliced transcript variants have been found. [provided by RefSeq, Aug 2020]</p> <p>Transcript Variant: This variant (7) lacks two alternate coding exons compared to variant 1, that causes a frameshift. This variant uses a downstream in-frame start-codon, so the encoded isoform 3 has a shorter N-terminus, as compared to isoform 1.</p>