

Product datasheet for SC332204

LCN1 (NM 001252618) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: LCN1 (NM_001252618) Human Untagged Clone

Tag: Tag Free Symbol: LCN1

Synonyms: PMFA; TLC; TP; VEGP

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC332204 representing NM_001252618.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GGTTCTCCATAA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001252618

Insert Size: 702 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001252618.1</u>

 RefSeq Size:
 785 bp

 RefSeq ORF:
 702 bp

 Locus ID:
 3933

 UniProt ID:
 P31025

 Cytogenetics:
 9q34.3

Protein Families: Secreted Protein

MW: 25.4 kDa

Gene Summary: This gene encodes a member of the lipocalin family of small secretory proteins. Lipocalins are

extracellular transport proteins that bind to a variety of hydrophobic ligands. The encoded protein is the primary lipid binding protein in tears and is overproduced in response to multiple stimuli including infection and stress. The encoded protein may be a marker for chromosome aneuploidy as well as an autoantigen in Sjogren's syndrome. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and two pseudogenes of this gene are also located on the long arm of chromosome 9. [provided

by RefSeq, Nov 2011]

Transcript Variant: This variant (3) uses an alternate splice site in the 3' coding region, which results in a translational frameshift, compared to variant 1. The encoded isoform (2) is longer

and has a distinct C-terminus, compared to isoform 1.