

## Product datasheet for **SC332643**

### NFAT1 (NFATC2) (NM\_001258297) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	NFAT1 (NFATC2) (NM_001258297) Human Untagged Clone
Tag:	Tag Free
Symbol:	NFAT1
Synonyms:	NFAT1; NFATP
Vector:	pCMV6-Entry (PS100001)



[View online »](#)

**Fully Sequenced ORF:** >SC332643 representing NM\_001258297.  
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

ATGTCACCTCGAACCAGCCTCGCCGAGGACAGCTGCCTGGGCCGCCACTCGCCCGTGCCCGTCCGGCC
TCCCGCTCCTCATCGCCTGGTGCCAAGCGGAGGCATTGTCGCCGAGGCCTTGGTTGCCCTGCCGCC
GGAGCCTCACCCAGCGCTCCCGAGCCCTCGCCGAGCCCTCATCTCACGTGGCACCCAGGACCAC
GGCTCCCGGCTGGGTACCCCTGTGGCTGGCTGCTGCCGTGATCATGGATGCCCTGAACAGCCTCGCC
ACGGACTCGCCTTGTGGGATCCCCCAAGATGTGGAAGACCAGCCCTGACCCCTCGCCGGTGTCTGCC
GCCCCATCCAAGGCCGGCTGCCTCGCCACATCTACCCGCGCTGGAGTTCTGGGGCCCTGCGAGCAG
GGCGAGAGGAGAACTCGGCTCCAGAATCCATCCTGCTGGTTCCGCCCACTTGGCCCAAGCCGCTGGTG
CCTGCCATTCCCATCTGCAGCATCCAGTGACTGCATCCCTCCCTCCACTTGAGTGGCCGCTGTCCAGT
CAGTCAGGCTCTTACGAGCTGCGGATCGAGGTGCAGCCCAAGCCACATCACCGGGCCACTATGAGACA
GAAGGCAGCCGAGGGGCTGTCAAAGCTCCAAGTGGAGGCCACCCTGTGGTTCAGTCCATGGCTACATG
GAAAACAAGCCTCTGGGACTTCAGATCTTATTGGGACAGCTGATGAGCGGATCCTTAAGCCGCACGCC
TTCTACCAGGTGCACCGAATCACGGGAAAAGTGTACCACCACCAGCTATGAGAAGATAGTGGGCAAC
ACCAAAGTCTGGAGATACCCCTTGAGGCCAAAAACAACATGAGGGCAACCATCGACTGTGCGGGGATC
TTGAAGCTTAGAAACGCCGACATTGAGCTGCGGAAAGGCGAGACGGACATTGGAAGAAAGAACCGCGG
GTGAGACTGGTTTTCCGAGTTACATCCCAGAGTCCAGTGGCAGAATCGTCTCTTTACAGACTGCATCT
AACCCCATCGAGTGCTCCAGCGATCTGCTCACGAGTGCCCATGGTTGAAAGACAAGACACAGACAGC
TGCTTGTCTATGGCGGCCAGCAATGATCCTCACGGGGCAGAACTTTACATCCGAGTCCAAAGTTGTG
TTTACTGAGAAGACCACAGATGGACAGCAAATTTGGGAGATGGAAGCCACGGTGGATAAGGACAAGAGC
CAGCCCAACATGCTTTTTTGTGAGATCCCTGAATATCGGAACAAGCATATCCGCACACCTGTAAGAGTG
AACTTCTACGTCATCAATGGGAAGAGAAAACGAAGTCAGCCTCAGCACTTTACCTACCACCCAGTCCCA
GCCATCAAGACGGAGCCACGGATGAATATGACCCCACTCTGATCTGCAGCCCCACCCATGGAGGCCTG
GGGAGCCAGCCTTACTACCCCGCAGCCCGATGGTGGCCGAGTCCCTCCTGCCTCGTGGCCACCATG
GCTCCCTGCCAGCAGTTCGCGACGGGGCTCTATCCCCTGACGCCGCTACCAGCAACAGAACCCAGCG
GCCGTACTTACCAGCGGAGCAAGAGCCTGAGCCCGAGCCTGCTGGGCTATCAGCAGCCGGCCCTCATG
GCCGCCCGCTGTCCCTTGGGACGCTCACCGCTCTGTGCTGGTGCAGCCGGCTCCAGGGCCAGAGC
TCAGCCCTGCTCCACCCCTCTCCGACCAACCAGCAGGCCCTCGCCTGTGATCCACTACTCACCCACCAAC
CAGCAGCTGCGCTGCGGAAGCCACCAGGAGTTCAGCACATCATGTACTGCGAGAATTTGCGACCAGGC
ACCACCAGACCTGGCCCGCCCGGTGAGTCAAGGTGAGAGGCTGAGCCCGGTTCTACCCACAGTC
ATTCAGCAGCAGAATGCCACGAGCCAAAGAGCCGCAAAAACGGACCCCGGTGAGTACCAAAAAGGAA
GTATTACCTGCGGGGTGACCATTAACAGGAGCAGAAGTGGACCAGACCTACTTGATGATGTTAAT
GAAATTATCAGGAAGGAGTTTTTCAGGACCTCTGCCAGAAATCAGACGTAA
  
```

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_001258297

**Insert Size:** 2121 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001258297.1</a>
<b>RefSeq Size:</b>	6928 bp
<b>RefSeq ORF:</b>	2121 bp
<b>Locus ID:</b>	4773
<b>Cytogenetics:</b>	20q13.2
<b>Protein Families:</b>	Druggable Genome, Transcription Factors
<b>Protein Pathways:</b>	Axon guidance, B cell receptor signaling pathway, MAPK signaling pathway, Natural killer cell mediated cytotoxicity, T cell receptor signaling pathway, VEGF signaling pathway, Wnt signaling pathway
<b>MW:</b>	77.1 kDa
<b>Gene Summary:</b>	<p>This gene is a member of the nuclear factor of activated T cells (NFAT) family. The product of this gene is a DNA-binding protein with a REL-homology region (RHR) and an NFAT-homology region (NHR). This protein is present in the cytosol and only translocates to the nucleus upon T cell receptor (TCR) stimulation, where it becomes a member of the nuclear factors of activated T cells transcription complex. This complex plays a central role in inducing gene transcription during the immune response. Alternate transcriptional splice variants encoding different isoforms have been characterized. [provided by RefSeq, Apr 2012]</p> <p>Transcript Variant: This variant (8) uses an alternate splice junction at the 5' end of a coding exon compared to variant 2. The resulting isoform (G) is shorter at the N-terminus compared to isoform C. Variants 6 and 8 both encode the same isoform (G). Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p>