

Product datasheet for SC332712

BLNK (NM 001258442) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: BLNK (NM_001258442) Human Untagged Clone

Tag: Tag Free Symbol: BLNK

Synonyms: AGM4; BASH; bca; BLNK-S; LY57; SLP-65; SLP65

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC332712 representing NM_001258442.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

ATGGACAAGCTTAATAAAATAACCGTCCCCGCCAGTCAGAAGTTGAGGCAGCTTCAAAAGATGGTCCAT
GATATTAAAAACAATGAAGGTGGAATAATGAATAAAATCAAAAAGCTAAAAGTCAAAAGCACCTCCAAGT
GTTCCTCGAAGGGACTACGCTTCAGAGAGCCCTGCTGACGAAGAGAGCAGTGGTCCGATGACTTTGAC
AGCGACTATGAAAATCCAGATGAGCACTCGGACTCAGAGATGTACGTGATGCCCGCCGAGGAGAACCGT
GATGACAGCTACGAGCCGCCTCCAGTAGAGCAGGAAACCAGGCCGGTTCACCCAGCCCTGCCCTTCGCC
AGAGGAACAGCTTCAGGTCGAAACAGTGGGGCCTGGGAAACCAAGTCACCTCCACCAGCTGCACCATCC
CCGTTGCCACGGGCCGGGAAAAAACCAACGACACCACTGAAGACAACTCCAGCTGCACCAGCTC
CCGTTGCCACGGGCCGGGAAAAAACCCATACCTGCACAGAAACCCACCGAGGGTCAAGTCACCAGACAAGAA
GCTGTCAGTTTTGTGAAGAAAAACCTATACCTGCTGAACACCACCACAAAAACCCATACCTCTGCCAAGA
TTTACAGAAGGGGGAAACCCAACTGTGGATGGCCCCTACCCAGCTTTTCATCTAATTCCACTATTTCA
GAACAGGAAGCTGGCGTTCTCTGCAAGCCATGGTATGCTGGAAGCCTGTGATCGAAAGTCTTCAACAAAAGCCTTTGCAACAAAACCCTTCAACACATAGTCCT
TTGGTTCTTATTGACAGTCAGAATAACACAAAAGATTCCACCAGACTGAAGTATGCAGTTAAAGTTTCA

 TAA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001258442

Insert Size: 900 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 001258442.1</u>

RefSeq Size: 1298 bp
RefSeq ORF: 900 bp
Locus ID: 29760
Cytogenetics: 10q24.1

Protein Families: Druggable Genome

Protein Pathways: B cell receptor signaling pathway, Primary immunodeficiency

MW: 33 kDa

Gene Summary: This gene encodes a cytoplasmic linker or adaptor protein that plays a critical role in B cell

development. This protein bridges B cell receptor-associated kinase activation with downstream signaling pathways, thereby affecting various biological functions. The phosphorylation of five tyrosine residues is necessary for this protein to nucleate distinct signaling effectors following B cell receptor activation. Mutations in this gene cause

hypoglobulinemia and absent B cells, a disease in which the pro- to pre-B-cell transition is developmentally blocked. Deficiency in this protein has also been shown in some cases of pre-B acute lymphoblastic leukemia. Alternatively spliced transcript variants have been found

Transcript Variant: This variant (5) lacks 3 in-frame exons in the mid- and 3' coding regions, and uses alternate in-frame donor and acceptor splice sites compared to variant 1, resulting

in a shorter isoform (5) compared to isoform 1.

for this gene. [provided by RefSeq, May 2012]