

## Product datasheet for SC332860

### PRC1 (NM\_001267580) Human Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** PRC1 (NM\_001267580) Human Untagged Clone  
**Tag:** Tag Free  
**Symbol:** PRC1  
**Synonyms:** ASE1  
**Vector:** pCMV6-Entry (PS100001)  
**Fully Sequenced ORF:** >SC332860 representing NM\_001267580.  
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

ATGAGGAGAAGTGAGGTGCTGGCGGAGGAGTCCATAGTATGTCTGCAGAAAGCCCTAAATCACCTTCGG
GAAATATGGGAGCTAATTGGGATTCAGAGGACCAGCGTTACAAAGAACTGAGGTGGTAAAGAAGCAT
ATCAAGGAAGAAGGAGAGACGACCATCTTGAACCTAGAAAAAGATTTGCGCACCCAAGTGGAAATGATG
CGAAAACAGAAAAGGAGAGAAAACAGGAACTGAAGCTACTTCAAGAGCAAGATCAAGAACTGTGCGAA
ATTCTTTGTATGCCCACTATGATATTGACAGTGCCTCAGTGCCCACTTGAAGAGCTGAACCACTTC
AGGCAACATGTGACAACCTTTGAGGGAAAACAAGGCTTCTAGGCGTGAGGAGTTTGTGAGTATAAAGAGA
CAGATCATACTGTGTATGGAAGAATTAGACCACACCCAGACACAAGCTTTGAAAGAGATGTGGTGTGT
GAAGACGAAGATGCCTTTTGTGCTTTGGAGAATATTGCAACACTACAAAAGTTGCTACGGCAGCTG
GAAATGCAGAAATCACAAAATGAAGCAGTGTGTGAGGGGCTGCGTACTCAAATCCGAGAGCTCTGGGAC
AGGTTGCAAAATACCTGAAGAAGAAAGAGAAGCTGTGGCCACCATTTATGTCTGGGTCAAAGGCCAAGGTC
CGGAAAGCGTGCAATTAGAAGTGGATCGGTTGGAAGAACTGAAAATGCAAAACATGAAGAAAGTGATT
GAGGCAATTCGAGTGGAGCTGGTTCAGTACTGGGACCAGTGTCTTTATAGCCAGGAGCAGAGACAAGCT
TTTGCCCTTTCTGTGCTGAGGACTACACAGAAAGTCTGCTCCAGCTCCACGATGCTGAGATTGTGCGG
TTAAAAAATACTATGAAGTTCACAAGGAACTCTTTGAAGTGTCCAGAAGTGGGAAAGAACTGGAGG
CTTTTCTTAGAGTTTGAGAGAAAAGCTTCAGATCCAAATCGATTTACAAACCGAGGAGGAAATCTTCTA
AAAGAAGAAAAACAACGAGCCAAGCTCCAGAAAATGCTGCCCAAGCTGGAAGAAGAGTTGAAGGCACGA
ATTGAATTGTGGGAACAGGAACATTCAAAGGCATTTATGGTGAATGGGCAGAAATTCATGGAGTATGTG
GCAGAACAATGGGAGATGCATCGATTGGAGAAAGAGAGCCAAGCAGGAAAGACAACCTGAAGAACAAA
AAACAGACAGAGACAGAGATGCTGTATGGCAGCGCTCCTCGAACACCTAGCAAGCGGCGAGGACTGGCT
CCCAATACACGGGCAAAGCACGTAAGCTGAACACTACCACCATGTCCAATGCTACGGCCAATAGTAGC
ATTCGGCCTATCTTTGGAGGGACAGTCTACCACCTCCCCGTGTCTCGACTTCTCCTTCTGGCAGCAAG
CCAGTCGCTGCTCCACCTGTTCAGGGAAGAAAACACCCCGTACTGGCAGGCATGGAGCCAACAAGGAG
AACCTGGAGCTCAACGGCAGCATCCTGAGTGCAGAACTTTCAAAGGCTTCCAAATCTGA
  
```

**Restriction Sites:** SgfI-MluI  
**ACCN:** NM\_001267580  
**Insert Size:** 1578 bp



[View online »](#)

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u>NM_001267580.1</u>
<b>RefSeq Size:</b>	2965 bp
<b>RefSeq ORF:</b>	1578 bp
<b>Locus ID:</b>	9055
<b>UniProt ID:</b>	<u>O43663</u>
<b>Cytogenetics:</b>	15q26.1
<b>MW:</b>	61.4 kDa
<b>Gene Summary:</b>	<p>This gene encodes a protein that is involved in cytokinesis. The protein is present at high levels during the S and G2/M phases of mitosis but its levels drop dramatically when the cell exits mitosis and enters the G1 phase. It is located in the nucleus during interphase, becomes associated with mitotic spindles in a highly dynamic manner during mitosis, and localizes to the cell mid-body during cytokinesis. This protein has been shown to be a substrate of several cyclin-dependent kinases (CDKs). It is necessary for polarizing parallel microtubules and concentrating the factors responsible for contractile ring assembly. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jun 2012]</p> <p>Transcript Variant: This variant (4) lacks an alternate in-frame exon in the 5' coding region, and uses an alternate splice site and lacks an exon that results in a frameshift in the 3' coding region, compared to variant 1. The encoded isoform (4) has a distinct C-terminus and is shorter than isoform 1.</p>