

Product datasheet for **SC332896**

BORIS (CTCFL) (NM_001269042) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	BORIS (CTCFL) (NM_001269042) Human Untagged Clone
Tag:	Tag Free
Symbol:	BORIS
Synonyms:	BORIS; CT27; CTCF-T; dj579F20.2; HMGB1L1
Vector:	pCMV6-Entry (PS100001)



[View online »](#)

Fully Sequenced ORF: >SC332896 representing NM_001269042.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

ATGGCAGCCACTGAGATCTCTGTCCTTTCTGAGCAATTCACCAAGATCAAAGAACTCGAGTTGATGCCG
GAAAAAGGCCCTGAAGGAGGAGGAAAAAGACGGAGTGTGCAGAGAGAAAGACCATCGGAGCCCTAGTGAG
TTGGAGGCCGAGCGTACCTCTGGGGCCTCCAGGACAGCGTCTGGAGGAAGAAGTGGAGCTGGTGCTG
GCCCCCTCGGAGGAGAGCGAGAAGTACATCCTGACCCTGCAGACGGTGCACCTTCACTTCTGAAGCTGTG
GAGTTGCAGGATATGAGCTTGTGAGCATACAGCAGCAAGAAGGGGTGCAGGTGGTGGTGAACAGCCT
GGCCCTGGGTTGCTGTGGCTTGAGGAAGGGCCCGGCAGAGCCTGCAGCAGTGTGTGGCCATTAGTATC
CAGCAAGAGCTGTACTCCCCGAAGAGATGGAGGTGTTGCAGTTCACGCTCTAGAGGAGAATGTGATG
GTGGCCAGTGAAGACAGTAAGTTAGCGGTGAGCCTGGCTGAAACTACTGGACTGATCAAGCTCGAGGAA
GAGCAGGAGAAGAACCAGTTATTGGCTGAAAGAACAAGGAGCAGCTCTTTTTTGTGAAACAATGTCA
GGAGATGAAAGAAGTGACGAAATGTTTCTCACAGTTTCAAATTCAAATGTGAAGAACAAGAGGATCAA
CCTACAGCTGGTCAAGCAGATGTGAAAAGGCCAAATCTACAAAAATCAAAGAAAGACAAAGGGAGCA
AAAGGAACCTTCCACTGTGATGTCTGCATGTTACCTTCTTAGAATGTCAAGTTTTAATCGTCATATG
AAAACACACACAGTGAAGCCTCACCTGTGTACCTCTGCCTGAAAACCTTCGTACGGTCACTCTG
CTGCGGAACCATGTTAACACCCACACAGGAACAGGCCCTACAAGTGTAACTGCAACATGGCATT
GTCACCACTGGAGAACTCGTCCGACACAGGCGCTATAACATACTCATGAGAAACCTTTAAATGTTCC
ATGTGCAAGTATGCCAGTGTGGAGGCAAGTAAATTGAAGCGCCATGTCCGATCCCACACTGGGGAGCGC
CCCTTTCAGTGTGCCAGTGCAGCTATGCCAGCAGAGATACCTACAAGCTGAAACGCCACATGAGAACG
CACTCAGGTGAGAAGCCTTACGAATGCCACATCTGCCACACCCGCTTACCCAGAGCGGGACCATGAAA
ATACATATTCTGCAGAAACACGGCGAAAATGTCCCAAATACCAGTGTCCCATTTGTGCCACCATCATT
GCACGAAAAAGCGACCTACGTGTGCATATGCCAACTTGCATGCTTACAGCGCTGCAGAGTAAAATGC
CGTACTGTTCTGCTGTCTTCCATGAACGCTATGCCCTCATTACGACCCAGAAAACCTATAAGAATGAG
AAGAGGTTCAAGTGCAAAACTGCAGTTATGCCTGCAAGCAGGAACGTCATATGACCGCTCACATTCGT
ACCCCACTGGAGAGAAACCATTCACCTGCCTTTCTTGAATAAATGTTCCGACAGAAGCAACTTCTA
AACGCTCACTTCAGGAAATACCACGATGCAAAATTCATCCGACTGTTTACAAATGCTCCAAGTGTGGC
AAAGGCTTTTCCGCTGGATTAACCTGCACAGACATTCGGAGAAGTGTGGATCAGGGGAAGCAAAGTCG
GCTGCTTCAGGAAAGGGAAGAAGAACAAGAAAGAGGAAGCAGACCATCTGAAGGAAGCCACAAAGGGT
CAGAAGGAAGCTGCGAAGGGATGGAAGGAAGCCGCAACGGAGACGCTGCTGCTGAGGAGGCTTCCACC
ACGAAGGGAGAACAGTTCAGGAGAGATGTTTCTGTGCCTGCAGAGAAACCACAGCCAGAGTCAAA
GAGGAAGTGGATGAAGGCGTGACCTGTGAAATGCTCCTCAACACGATGGATAAGTGA
  
```

Restriction Sites: SgfI-MluI

ACCN: NM_001269042

Insert Size: 1989 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001269042.1](#)

RefSeq Size: 3598 bp

RefSeq ORF: 1989 bp

Locus ID: 140690

UniProt ID: [Q8NI51](#)

Cytogenetics: 20q13.31

Protein Families: Transcription Factors

MW: 75.6 kDa

Gene Summary: CCCTC-binding factor (CTCF), an 11-zinc-finger factor involved in gene regulation, utilizes different zinc fingers to bind varying DNA target sites. CTCF forms methylation-sensitive insulators that regulate X-chromosome inactivation. This gene is a paralog of CTCF and appears to be expressed primarily in the cytoplasm of spermatocytes, unlike CTCF which is expressed primarily in the nucleus of somatic cells. CTCF and the protein encoded by this gene are normally expressed in a mutually exclusive pattern that correlates with resetting of methylation marks during male germ cell differentiation. Multiple alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jun 2012]

Transcript Variant: This variant (4) has an alternate splice site in the 3' coding region, compared to variant 1. The resulting isoform (2) lacks an internal aa in the C-terminal region, compared to isoform 1.