

Product datasheet for **SC332944**

WWP2 (NM_001270453) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	WWP2 (NM_001270453) Human Untagged Clone
Tag:	Tag Free
Symbol:	WWP2
Synonyms:	AIP2; WWp2-like
Vector:	pCMV6-Entry (PS100001)



[View online »](#)

Fully Sequenced ORF: >SC332944 representing NM_001270453.
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

```

ATGCAGCTGACCCTGAACCTGCAGACGGAGAACAAGGCAGCGTTGTCTCAGGCGGAGAGCTGACAATT
TTCCTGGACGGGCAACTGTTGATCTGGGAAATGTGCCTAATGGCAGTGCCCTGACAGATGGATCACAG
CTGCCTTCGAGAGACTCCAGTGGAACAGCAGTAGCTCCAGAGAACCGGCACCAGCCCCAGCACAAC
TGCTTTGGTGAAGATCCGGACGCACAGACATTGGGTGCTTCAGCCAGAACAACCCAGCAACCGGC
GAGCAAAGCCCCGGTGTCTCGGAGCCGGCACCGCCAGCCCGTCAAGAACTCAGGCCACAGTGGCTTGGCC
AATGGCACAGTGAATGATGAACCCACAACAGCCACTGATCCCGAAGAACCCTCCGTTGTTGGTGTGACG
TCCCCACCTGCTGACCCCTTGAGTGTGACCCCGAATCCCAACACGACTTCTCTCCCTGCCCCAGCCACA
CCGGCTGAAGGAGAGGAACCCAGCACTTCGGGTACACAGCAGCTCCAGCGGCTGCCAGGCCCCCGAC
GCTCTGCCTGCTGGATGGGAACAGCAGAGCTGCCAACGGACGTGTCTATTATGTTGACCACAATACC
AAGACCACCACCTGGGAGCGGCCCTTCTCCAGGCTGGGAAAAACGCACAGATCCCGAGGCAGGTTT
TACTATGTGGATCACAATACTCGGACCACCACCTGGCAGCGTCCGACCGGGAGTACGTGCGCAACTAT
GAGCAGTGGCAGTCGACGGGAATCAGTCCAGGGGCCATGCAGCACTTCAGCCAAAGATTCTCTAC
CAGTCTTCGAGTCTTCGACTGACCATGATCCCTGGGCCCTCCCTCTGGCTGGGAGAAGAGACAG
GACAATGGACGGGTGATTACGTGAACCATAAACTCGCACGACCCAGTGGGAGGATCCCGGACCCAG
GGGATGATCCAGGAACAGCTCTGCCCCAGGATGGGAGATGAAATACACCAGCGAGGGGGTGGCATA
TTTGTGGACCACAATACCCGACCCACCCTTTAAGGATCCTCGCCCGGGTTTGGTGGGACGAAG
CAAGGTTCCCTGGTGTATGACCGCAGTTTTCGGTGGAAGTATCACCAGTTCGGTTTCTCTGCCAT
TCAAATGCCCTACCTAGCCACGTGAAGTACAGCGTTTCCAGGCAGACGTTTTCGAAGATTCTTCCAA
CAGATCATGAACATGAAACCCTATGACCTGCGCCGCGGCTCTACATCATATGCGTGGCAGGAGGGC
CTGGACTATGGGGCATCGCCAGAGAGTGGTTTTTCTCTGTCTCATGAGGTGCTCAACCCTATGTAT
TGTTTTATTGAATATGCCGGAAGAACAATTACTGCCTGCAGATCAACCCGCTCTCCATCAACCCG
GACCACCTCACCTACTTTCGTTTTATAGGCAGATTTCATGCCATGGCGCTGTACCATGGAAGTTTCATC
GACACGGGCTTACCCTCCCTTTCTACAAGCGGATGCTCAATAAGAGACCAACCTGAAAGACTGGAG
TCCATTGACCCTGAGTTCTACAACCTCATTGTCTGGATCAAAGAGAACAACCTGGAAGAATGTGGCTG
GAGCTGTACTTCCAGGACATGGAGATACTGGGCAAGGTGACGACCCACGAGCTGAAGGAGGGCGGC
GAGAGCATCCGGTCCAGAGGAGAACAAGGAAGGTACATCATGCTGCTGACTGACTGGCGTTTACC
CGAGGCGTGAAGAGCAGACCAAAGCCTTCTGGATGGCTTCAACGAGGTGGCCCCGCTGGAGTGGCTG
CGTACTTTGACGAGAAAGAGCTGGAGCTGATGCTGTGCGGCATGCAGGAGATAGACATGAGCGACTGG
CAGAAGAGCACCATCTACCGGCACTACACCAAGAACAGCAAGCAGATCCAGTGGTCTGGCAGGTGGTG
AAGGAGATGGACAACGAGAAGAGGATCCGGCTGCTGCAGTTTGTACCGGTACCTGCCGCTGCCGCTC
GGGGGATTTGCCGAACCTCATCGGTAGCAACGGACCACAGAAGTTTTGCATTGACAAAGTTGGCAAGGAA
ACCTGGCTGCCGAGAAGCCACACCTGCTTCAACCGTCTGGATCTTCCACCCTACAAGAGCTACGAACAG
CTGAGAGAGAAGCTGCTGTATGCCATTGAGGAGACCGAGGGCTTTGGACAGGAGTAA
  
```

Restriction Sites: SgfI-MluI

ACCN: NM_001270453

Insert Size: 2265 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001270453.1](#)

RefSeq Size: 4245 bp

RefSeq ORF: 2265 bp

Locus ID: 11060

UniProt ID: [O00308](#)

Cytogenetics: 16q22.1

Protein Families: Druggable Genome

Protein Pathways: Ubiquitin mediated proteolysis

MW: 86.3 kDa

Gene Summary: This gene encodes a member of the Nedd4 family of E3 ligases, which play an important role in protein ubiquitination. The encoded protein contains four WW domains and may play a role in multiple processes including chondrogenesis and the regulation of oncogenic signaling pathways via interactions with Smad proteins and the tumor suppressor PTEN. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 10. [provided by RefSeq, Jul 2012]

Transcript Variant: This variant (4) differs in the 5' UTR, lacks a portion of the 5' coding region and initiates translation at a downstream, in-frame start codon, compared to variant 1. The encoded isoform (4) has a shorter N-terminus, compared to isoform 1.