

Product datasheet for SC333093

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SPINK2 (NM_001271720) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: SPINK2 (NM_001271720) Human Untagged Clone

Tag: Tag Free Symbol: SPINK2

Synonyms: HUSI-II; SPGF29

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC333093 representing NM_001271720.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

AATGGACCCTGCTGA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001271720

Insert Size: 360 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.



SPINK2 (NM_001271720) Human Untagged Clone - SC333093

RefSeq: <u>NM 001271720.1</u>

RefSeq Size: 741 bp
RefSeq ORF: 360 bp
Locus ID: 6691
Cytogenetics: 4q12

Protein Families: Secreted Protein, Transmembrane

MW: 12.5 kDa

Gene Summary: This gene encodes a member of the family of serine protease inhibitors of the Kazal type

(SPINK). The encoded protein acts as a trypsin and acrosin inhibitor in the genital tract and is localized in the spermatozoa. The protein has been associated with the progression of lymphomas. Alternative splicing results in multiple transcript variants. [provided by RefSeq,

Nov 2012]

Transcript Variant: This variant (4) uses an alternate in-frame splice site in the 5' coding region, compared to variant 1. It encodes isoform 4 which is shorter than isoform 1.