

## **Product datasheet for SC333164**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## C18orf10 (TPGS2) (NM\_001271950) Human Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

Product Name: C18orf10 (TPGS2) (NM\_001271950) Human Untagged Clone

Tag: Tag Free
Symbol: C18orf10

Synonyms: C18orf10; HMFN0601; L17; PGs2

**Vector:** pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC333164 representing NM\_001271950.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

Restriction Sites: Sgfl-Mlul

**ACCN:** NM\_001271950

**Insert Size:** 774 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** NM 001271950.1

 RefSeq Size:
 3939 bp

 RefSeq ORF:
 774 bp

 Locus ID:
 25941

 UniProt ID:
 Q68CL5

 Cytogenetics:
 18q12.2

 MW:
 28.7 kDa

**Gene Summary:** This gene encodes a protein that is a component of the neuronal polyglutamylase complex,

which plays a role in post-translational addition of glutamate residues to C-terminal tubulin tails. Alternatively spliced transcript variants encoding multiple isoforms have been observed

for this gene. [provided by RefSeq, Dec 2012]

Transcript Variant: This variant (3) lacks an alternate in-frame exon in the central coding region compared to variant 1. It encodes isoform 3 which is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used

for the transcript record were based on transcript alignments.