

Product datasheet for SC333267

i roddet datasneet for sessser

IFTAP (NM_001276724) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: IFTAP (NM_001276724) Human Untagged Clone

Tag: Tag Free
Symbol: IFTAP

Synonyms: C11orf74; HEPIS; NWC

Vector: pCMV6-Entry (PS100001)

Fully Sequenced ORF: >SC333267 representing NM_001276724.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

AAGAGAAAGGACACCAGCCCAGACTTAGAGAAATCCTGTGAC<mark>TGA</mark>

Restriction Sites: Sgfl-Mlul

ACCN: NM 001276724

Insert Size: 666 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM 001276724.1

 RefSeq Size:
 955 bp

 RefSeq ORF:
 666 bp

 Locus ID:
 119710

 UniProt ID:
 Q86VG3

 Cytogenetics:
 11p12

 MW:
 25.4 kDa

Gene Summary: This gene encodes a protein that was identified as a cellular interacting partner of non-

structural protein 10 of the severe acute respiratory syndrome coronavirus (SARS-CoV). The encoded protein may function as a negative regulator of transcription. There is a pseudogene for this gene on chromosome 1. Alternative splicing results in multiple transcript variants.

[provided by RefSeg, Mar 2013]

Transcript Variant: This variant (3) differs in the 5' UTR, compared to variant 1. Variants 1, 2, 3,

and 4 encode the same isoform (a).