

## Product datasheet for **SC333381**

### Protamine 2 (PRM2) (NM\_001286357) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Protamine 2 (PRM2) (NM_001286357) Human Untagged Clone
Tag:	Tag Free
Symbol:	Protamine 2
Synonyms:	CT94.2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC333381 representing NM_001286357. Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGAATTCGTCGACTG  
 GATCCGGTACCGAGGAGATCTGCCGCC**CGATCGCC**  
 ATGGTCCGATACCGCGTGAGGAGCCTGAGCGAACGCTCGCACGAGGTGTACAGGCAGCAGTTGCATGGG  
 CAAGAGCAAGGACACCGGCCAAGAGGAGCAAGGGCTGAGCCCGGAGCACGTCGAGGTCTACGAGAGG  
 ACCCATGGCCAGTCTCACTATAGGCGCAGACACTGCTCTCGAAGGAGGCTGCACCGGATCCACAGCGG  
 CAGCATCGCTCCTGCAGAAGGCAC**TAA**  
**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT  
 TACAAGGATGACGACGATAAGGTTAAACGGCCGGC

Restriction Sites:	SgfI-MluI
ACCN:	NM_001286357
Insert Size:	234 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).


[View online »](#)

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001286357.1](#)

**RefSeq Size:** 608 bp

**RefSeq ORF:** 234 bp

**Locus ID:** 5620

**Cytogenetics:** 16p13.13

**MW:** 9.7 kDa

**Gene Summary:** Protamines substitute for histones in the chromatin of sperm during the haploid phase of spermatogenesis, and are the major DNA-binding proteins in the nucleus of sperm in many vertebrates. They package the sperm DNA into a highly condensed complex in a volume less than 5% of a somatic cell nucleus. Many mammalian species have only one protamine (protamine 1); however, a few species, including human and mouse, have two. This gene encodes protamine 2, which is cleaved to give rise to a family of protamine 2 peptides. Alternatively spliced transcript variants have also been found for this gene. [provided by RefSeq, Sep 2015]

Transcript Variant: This variant (3) uses alternate donor and acceptor splice sites at the junction of exons 1 and 2 compared to variant 1. The resulting isoform (3) lacks an internal protein segment and has a shorter C-terminus compared to isoform 1.