

Product datasheet for **SC333393**

UBE2C (NM_001281742) Human Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	UBE2C (NM_001281742) Human Untagged Clone
Tag:	Tag Free
Symbol:	UBE2C
Synonyms:	dj447F3.2; UBCH10
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC333393 representing NM_001281742. Blue=Insert sequence Red=Cloning site Green=Tag(s)

```
GCTCGTTTAGTGAACCGTCAGAATTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCCCGATCGCC
ATGGCTTCCAAAACCGGACCCAGCCGCACTAGCGTCGCCGCCCGCCGTAAGGAGCTGAGCCGAGC
GGGGCGCCCGCGGGTCCGGTGGCAAAGGCTACAGCAGGAGCTGATGACCCATGATGTCTGGC
GATAAAGGGATTCTGCCTTCCCTGAATCAGACAACCTTTCAAATGGGTAGGGACCATCCATGGAGCA
GCTGGAACAAACCAACATTGATAGTCCCTGA
ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC
```

Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001281742
Insert Size:	240 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001281742.1](#)

RefSeq Size: 659 bp

RefSeq ORF: 240 bp

Locus ID: 11065

UniProt ID: [O00762](#)

Cytogenetics: 20q13.12

Protein Families: Druggable Genome

Protein Pathways: Ubiquitin mediated proteolysis

MW: 8.1 kDa

Gene Summary: The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013]
Transcript Variant: This variant (8) lacks an exon in central coding region, which results in a translational frameshift, compared to variant 1. The encoded isoform (7) has a shorter and distinct C-terminus, compared to isoform 1.