

## Product datasheet for **SC333414**

### Apc11 (ANAPC11) (NM\_001289415) Human Untagged Clone


#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Apc11 (ANAPC11) (NM\_001289415) Human Untagged Clone  
**Tag:** Tag Free  
**Symbol:** ANAPC11  
**Synonyms:** APC11; Apc11p; HSPC214  
**Vector:** pCMV6-Entry (PS100001)  
**Fully Sequenced ORF:** >SC333414 representing NM\_001289415.  
 Blue=Insert sequence Red=Cloning site Green=Tag(s)

ATGAAGGTGAAGATTAAGTGCTGGAACGGCGTGGCCACTTGGCTCTGGGTGGCCAACGATGAGAAGTGT  
 GGCATCTGCAGGATGGCATTTAACGGATGCTGCCCTGACTGCAAGGTGCCCGGCGACGACTGCCCGCTG  
 GTGTGGGGCCAGTGTCCACTGCTTCCACATGCATTGCATCCTCAAGTGGCTGCACGCACAGCAGGTG  
 CAGCAGCACTGCCCCATGTGCCCCAGGAATGGAAGTTCAAGGAGTGA

**Restriction Sites:** SgfI-MluI

#### Plasmid Map:



**ACCN:** NM\_001289415

**Insert Size:** 255 bp



[View online >](#)

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_001289415.1</a>
<b>RefSeq Size:</b>	885 bp
<b>RefSeq ORF:</b>	255 bp
<b>Locus ID:</b>	51529
<b>UniProt ID:</b>	<a href="#">Q9NYG5</a>
<b>Cytogenetics:</b>	17q25.3
<b>Protein Families:</b>	Druggable Genome
<b>Protein Pathways:</b>	Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation, Ubiquitin mediated proteolysis
<b>MW:</b>	9.8 kDa
<b>Gene Summary:</b>	<p>Together with the cullin protein ANAPC2, constitutes the catalytic component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the cell cycle. The APC/C complex acts by mediating ubiquitination and subsequent degradation of target proteins: it mainly mediates the formation of 'Lys-11'-linked polyubiquitin chains and, to a lower extent, the formation of 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains. May recruit the E2 ubiquitin-conjugating enzymes to the complex.[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (9) lacks an exon in the 3' coding region, which results in a frameshift, compared to variant 1. The encoded isoform (2) has a distinct C-terminus and is shorter than isoform 1. Variants 2, 3, 4, 5, 6, 7, 8, 9, 10, and 11 encode the isoform 2.</p>