

## Product datasheet for **SC333537**

### Cyclophilin A (PPIA) (NM\_001300981) Human Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Cyclophilin A (PPIA) (NM_001300981) Human Untagged Clone
Tag:	Tag Free
Symbol:	Cyclophilin A
Synonyms:	CYPA; CYPH; HEL-S-69p
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>SC333537 representing NM_001300981. Blue=Insert sequence Red=Cloning site Green=Tag(s)

GCTCGTTTAGTGAACCGTCAGAATTTTGTAAACGACTCACTATAGGGCGGCCGGAATTCGTCGACTG  
 GATCCGGTACCGAGGAGATCTGCCGCC**CGATCGCC**  
 ATGTGTCAGGGTGGTGAATTCACACGCCATAATGGCACTGGTGGCAAGTCCATCTATGGGAGAAATTT  
 GAAGATGAGAATTCATCCTAAAGCATACGGTCCTGGCATCTTGCCATGGCAAATGCTGGACCAAC  
 ACAAATGGTTCCAGTTTTTCATCTGCACTGCCAAGACTGAGTGGTTGGATGGCAAGCATGTGGTGT  
 GGCAAAGTGAAAGAAGGCATGAATATTGTGGAGCCATGGAGCGCTTTGGGTCCAGGAATGGCAAGACC  
 AGCAAGAAGATCACCATTGCTGACTGTGGACAAC**TGAA**  
**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT  
 TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites:	SgfI-MluI
ACCN:	NM_001300981
Insert Size:	318 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).


[View online »](#)

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_001300981.1](#)

**RefSeq Size:** 2445 bp

**RefSeq ORF:** 318 bp

**Locus ID:** 5478

**UniProt ID:** [P62937](#)

**Cytogenetics:** 7p13

**MW:** 11.4 kDa

**Gene Summary:** This gene encodes a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. The encoded protein is a cyclosporin binding-protein and may play a role in cyclosporin A-mediated immunosuppression. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions. Multiple pseudogenes that map to different chromosomes have been reported. [provided by RefSeq, Jul 2008]  
 Transcript Variant: This variant (2) differs in its 5' UTR and uses a downstream start codon, compared to variant 1. The encoded isoform (2) has a shorter N-terminus, compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.