

Product datasheet for SC333599

POLR2H (NM 001278700) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: POLR2H (NM_001278700) Human Untagged Clone

Tag: Tag Free Symbol: POLR2H

Synonyms: RPABC3; RPB8; RPB17

Mammalian Cell

Selection:

Neomycin

Vector: pCMV6-Entry (PS100001) **E. coli Selection:** Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC333599 representing NM_001278700.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001278700

Insert Size: 345 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001278700.1</u>



POLR2H (NM_001278700) Human Untagged Clone - SC333599

RefSeq Size: 1114 bp
RefSeq ORF: 345 bp
Locus ID: 5437
UniProt ID: P52434
Cytogenetics: 3q27.1

Protein Families: Transcription Factors

Protein Pathways: Huntington's disease, Metabolic pathways, Purine metabolism, Pyrimidine metabolism, RNA

polymerase

MW: 13 kDa

Gene Summary: The three eukaryotic RNA polymerases are complex multisubunit enzymes that play a central

role in the transcription of nuclear genes. This gene encodes an essential and highly conserved subunit of RNA polymerase II that is shared by the other two eukaryotic DNA-directed RNA polymerases, I and III. Alternative splicing results in multiple transcript variants

of this gene. [provided by RefSeq, Jul 2013]

Transcript Variant: This variant (4) contains an alternate 5' terminal exon, initiates translation

at a downstream start codon and uses an alternate splice site in the 3' coding region, resulting in a frameshift, compared to variant 1. Variants 3 and 4 encode the same isoform

(3), which is shorter and has a distinct C-terminus, compared to isoform 1.