

Product datasheet for SC333625

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MDMX (MDM4) (NM_001278518) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: MDMX (MDM4) (NM_001278518) Human Untagged Clone

Tag: Tag Free Symbol: MDM4

Synonyms: BMFS6; HDMX; MDMX; MRP1

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC333625 representing NM_001278518.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

GATTGA


ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul



Plasmid Map:



ACCN: NM_001278518

Insert Size: 351 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001278518.1</u>



MDMX (MDM4) (NM_001278518) Human Untagged Clone - SC333625

RefSeq Size: 9611 bp
RefSeq ORF: 351 bp
Locus ID: 4194
UniProt ID: O15151
Cytogenetics: 1q32.1

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: p53 signaling pathway

MW: 12.8 kDa

Gene Summary: This gene encodes a nuclear protein that contains a p53 binding domain at the N-terminus

and a RING finger domain at the C-terminus, and shows structural similarity to p53-binding protein MDM2. Both proteins bind the p53 tumor suppressor protein and inhibit its activity, and have been shown to be overexpressed in a variety of human cancers. However, unlike MDM2 which degrades p53, this protein inhibits p53 by binding its transcriptional activation domain. This protein also interacts with MDM2 protein via the RING finger domain, and inhibits the latter's degradation. So this protein can reverse MDM2-targeted degradation of p53, while maintaining suppression of p53 transactivation and apoptotic functions.

Alternatively spliced transcript variants encoding different isoforms have been noted for this

gene. [provided by RefSeq, Feb 2011]

Transcript Variant: This variant (6, also known as MDM4-ALT1, XALT1, or HDMX-ALT1) lacks four consecutive exons compared to variant 1. The resulting isoform (6) has a shorter and distinct C-terminus compared to isoform 1. Annotation of this protein causes this transcript to be a candidate for nonsense-mediated mRNA decay (NMD); however, since transcript variant 4 escapes NMD, it is likely that isoform 6 is also biologically valid. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.