

Product datasheet for SC333645

MTLRP (GHRL) (NM_001302824) Human Untagged Clone

Product data:

Product Type: Expression Plasmids Product Name: MTLRP (GHRL) (NM_001302824) Human Untagged Clone Tag: Tag Free Symbol: GHRL MTLRP Synonyms: **Mammalian Cell** Neomycin Selection: Vector: pCMV6-Entry (PS100001) E. coli Selection: Kanamycin (25 ug/mL) >SC333645 representing NM_001302824. **Fully Sequenced ORF:** Blue=Insert sequence Red=Cloning site Green=Tag(s) GCTCGTTTAGTGAACCGTCAGAATTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTG GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC ATGCCCTCCCCAGGGACCGTCTGCAGCCTCCTGCTCCTCGGCATGCTCTGGCCGACTTGGCCATGGCA GGCTCCAGCTTCCTGAGCCCTGAACACCAGAGAGTCCAGCAGAGAAGGAGTCGAAGAAGCCACCAGCC AAGCTGCAGCCCCGAGCTCTAGCAGGCTGGCTCCGCCCGGAAGATGGAGGTCAAGCAGAAGGGGCAGAG GATGAACTGGAAGTCCGGTTCAACGCCCCCTTTGATGTTGGAATCAAGCTGTCAGGGGTTCAGTACCAG CAGCACAGCCAGGCCCTGGGGAAGTTTCTTCAGGACATCCTCTGGGAAGAGGCCCAAAGAGGCCCCAGCC GACAAGTGA ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites:

Sgfl-Mlul


OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Plasmid Map:



| ACCN: | NM_001302824 |
|------------------------|--|
| Insert Size: | 354 bp |
| OTI Disclaimer: | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP). |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM 001302824.1</u> |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | MTLRP (GHRL) (NM_001302824) Human Untagged Clone – SC333645 |
|--|---|
|--|---|

| RefSeq Size: | 835 bp |
|-------------------|--|
| RefSeq ORF: | 354 bp |
| Locus ID: | 51738 |
| UniProt ID: | Q9UBU3 |
| Cytogenetics: | 3p25.3 |
| Protein Families: | Druggable Genome, Secreted Protein, Transmembrane |
| MW: | 12.9 kDa |
| Gene Summary: | This gene encodes the ghrelin-obestatin preproprotein that is cleaved to yield two peptides, ghrelin and obestatin. Ghrelin is a powerful appetite stimulant and plays an important role in energy homeostasis. Its secretion is initiated when the stomach is empty, whereupon it binds to the growth hormone secretagogue receptor in the hypothalamus which results in the secretion of growth hormone (somatotropin). Ghrelin is thought to regulate multiple activities, including hunger, reward perception via the mesolimbic pathway, gastric acid secretion, gastrointestinal motility, and pancreatic glucose-stimulated insulin secretion. It was initially proposed that obestatin plays an opposing role to ghrelin by promoting satiety and thus decreasing food intake, but this action is still debated. Recent reports suggest multiple metabolic roles for obestatin, including regulating adipocyte function and glucose metabolism. Alternative splicing results in multiple transcript variants. In addition, antisense transcripts for this gene have been identified and may potentially regulate ghrelin-obestatin preproprotein expression. [provided by RefSeq, Nov 2014] Transcript Variant: This variant (11) differs in the 5' UTR compared to variant 1. This isoform (11) contains the ligands ghrelin-28 and obestatin. Variants 1, 8, 9, 11 and 12 encode the same protein. |

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US