

Product datasheet for SC333650

POLR2F (NM 001301129) Human Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: POLR2F (NM_001301129) Human Untagged Clone

Tag: Tag Free
Symbol: POLR2F

Synonyms: HRBP14.4; POLRF; RPABC2; RPABC14.4; RPB6; RPB14.4; RPC15

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >SC333650 representing NM_001301129.

Blue=Insert sequence Red=Cloning site Green=Tag(s)

GATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATCACCGACTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT

TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Plasmid Map:



ACCN: NM_001301129

Insert Size: 357 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001301129.1</u>



POLR2F (NM_001301129) Human Untagged Clone - SC333650

RefSeq Size: 2207 bp
RefSeq ORF: 357 bp
Locus ID: 5435
Cytogenetics: 22q13.1

Protein Families: Transcription Factors

Protein Pathways: Huntington's disease, Metabolic pathways, Purine metabolism, Pyrimidine metabolism, RNA

polymerase

MW: 13.3 kDa

Gene Summary: This gene encodes the sixth largest subunit of RNA polymerase II, the polymerase responsible

for synthesizing messenger RNA in eukaryotes. In yeast, this polymerase subunit, in combination with at least two other subunits, forms a structure that stabilizes the transcribing polymerase on the DNA template. Alternative splicing results in multiple

transcript variants. [provided by RefSeq, Jul 2014]

Transcript Variant: This variant (2) contains an alternate exon in the 5' region and initiates translation at an alternate start codon, compared to variant 1. The encoded isoform (2) has a distinct N-terminus and is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record

were based on transcript alignments.