

## Product datasheet for SC333657

### IFI27 (NM\_001288957) Human Untagged Clone


#### Product data:

Product Type:	Expression Plasmids
Product Name:	IFI27 (NM_001288957) Human Untagged Clone
Tag:	Tag Free
Symbol:	IFI27
Synonyms:	FAM14D; ISG12; ISG12A; P27
Vector:	pCMV6-Entry (PS100001)
Fully Sequenced ORF:	>SC333657 representing NM_001288957. Blue=Insert sequence Red=Cloning site Green=Tag(s)

ATGGAGGCTCTGCTCTCACCTCATCAGCAGTGACCAGTGTGGCCAAAGTGGTCAGGGTGGCCTCTGCC  
TCTGCCGTAGTTTTGCCCTGGCCAGGATTGCTACAGTTGTGATTGGAGGAGTTGTGGCTGTGCCCATG  
GTGCTCAGTGCCATGGGCTTCACTGCGGCGGAATCGCCTCGTCTCCATAGCAGCCAAGATGATGTCC  
GCGGCGGCCATTGCCAATGGGGTGGAGTTGCCTCGGCGAGCCTTGTGGCTACTCTGCAGTCACTGGGA  
GCAACTGGACTCTCCGATTGACCAAGTTCATCCTGGGCTCCATTGGGTCTGCCATTGCGGCTGTCATT  
GCGAGTTCTACTAG

Restriction Sites: SgfI-MluI

Plasmid Map:



ACCN: NM\_001288957



[View online >](#)

<b>Insert Size:</b>	360 bp
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_001288957.1</a></u>
<b>RefSeq Size:</b>	737 bp
<b>RefSeq ORF:</b>	360 bp
<b>Locus ID:</b>	3429
<b>UniProt ID:</b>	<u><a href="#">P40305</a></u>
<b>Cytogenetics:</b>	14q32.12
<b>Protein Families:</b>	Transmembrane
<b>MW:</b>	11.3 kDa

**Gene Summary:**

Probable adapter protein involved in different biological processes (PubMed:22427340, PubMed:27194766). Part of the signaling pathways that lead to apoptosis (PubMed:18330707, PubMed:27673746, PubMed:24970806). Involved in type-I interferon-induced apoptosis characterized by a rapid and robust release of cytochrome C from the mitochondria and activation of BAX and caspases 2, 3, 6, 8 and 9 (PubMed:18330707, PubMed:27673746). Also functions in TNFSF10-induced apoptosis (PubMed:24970806). May also have a function in the nucleus, where it may be involved in the interferon-induced negative regulation of the transcriptional activity of NR4A1, NR4A2 and NR4A3 through the enhancement of XPO1-mediated nuclear export of these nuclear receptors (PubMed:22427340). May thereby play a role in the vascular response to injury (By similarity). In the innate immune response, has an antiviral activity towards hepatitis C virus/HCV (PubMed:27194766, PubMed:27777077). May prevent the replication of the virus by recruiting both the hepatitis C virus non-structural protein 5A/NS5A and the ubiquitination machinery via SKP2, promoting the ubiquitin-mediated proteasomal degradation of NS5A (PubMed:27194766, PubMed:27777077). [UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (6) differs in the 5' UTR and lacks a 9-nt segment in the CDS, as compared to variant 1. The resulting isoform (2) lacks an internal 3 aa segment, as compared to isoform 1. Variants 2, 4, 6, 7 and 10 encode the same isoform.